

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number  
WO 01/87937 A2(51) International Patent Classification<sup>7</sup>: C07K 14/00

(21) International Application Number: PCT/US01/16285

(22) International Filing Date: 17 May 2001 (17.05.2001)

(25) Filing Language: English

(26) Publication Language: English

## (30) Priority Data:

60/205,628	18 May 2000 (18.05.2000)	US
60/206,222	22 May 2000 (22.05.2000)	US
60/207,566	25 May 2000 (25.05.2000)	US
60/208,834	2 June 2000 (02.06.2000)	US
60/208,861	2 June 2000 (02.06.2000)	US

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PATTERSON, Chandra [US/US]; 490 Sherwood Way, #1, Menlo Park, CA 94025 (US). LU, Dyung, Aina, M. [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). THORNTON, Michael [US/US]; 9 Medway Road, Woodside, CA 94062-2612 (US). LU, Yan [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). TRIBOULEY, Catherine, M. [FR/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). GRAUL, Richard [US/US]; 682-29th Avenue, #104, San Francisco, CA 94121 (US). KHAN, Farrah, A. [IN/US]; 333 Escuela Avenue, #221, Mountain View, CA 94040 (US). GANDHI, Ammeena, R. [US/US]; 837 Roble Avenue, #1, Menlo Park, CA 94025 (US). WALIA, Narinder, K. [US/US]; 890 Davis Street, #205, San Leandro, CA 94577 (US). NGUYEN, Danniell, B. [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118

(US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). HAFALIA, April [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). ELLIOTT, Vicki, S. [US/US]; 3770 Poltan Place Way, San Jose, CA 95121 (US). LAL, Preeti [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). REDDY, Roopa [IN/US]; 1233 W. McKinley Avenue, #3, Sunnyvale, CA 94086 (US). KALLICK, Deborah, A. [US/US]; 900 Olive Street, Menlo Park, CA 94025 (US). TANG, Tom, Y. [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US).

(74) Agents: HAMLET-COX, Diana, et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: G-PROTEIN COUPLED RECEPTORS

(57) Abstract: The invention provides human G-protein coupled receptors (GCREC) and polynucleotides which identify and encode GCREC. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of GCREC.

WO 01/87937 A2

## G-PROTEIN COUPLED RECEPTORS

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of G-protein coupled  
5 receptors and to the use of these sequences in the diagnosis, treatment, and prevention of cell  
proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic  
disorders, and viral infections, and in the assessment of the effects of exogenous compounds on the  
expression of nucleic acid and amino acid sequences of G-protein coupled receptors.

### BACKGROUND OF THE INVENTION

Signal transduction is the general process by which cells respond to extracellular signals.  
Signal transduction across the plasma membrane begins with the binding of a signal molecule, e.g., a  
hormone, neurotransmitter, or growth factor, to a cell membrane receptor. The receptor, thus  
activated, triggers an intracellular biochemical cascade that ends with the activation of an intracellular  
15 target molecule, such as a transcription factor. This process of signal transduction regulates all types  
of cell functions including cell proliferation, differentiation, and gene transcription. The G-protein  
coupled receptors (GPCRs), encoded by one of the largest families of genes yet identified, play a  
central role in the transduction of extracellular signals across the plasma membrane. GPCRs have a  
proven history of being successful therapeutic targets.

20 GPCRs are integral membrane proteins characterized by the presence of seven hydrophobic  
transmembrane domains which together form a bundle of antiparallel alpha ( $\alpha$ ) helices. GPCRs range  
in size from under 400 to over 1000 amino acids (Strosberg, A.D. (1991) Eur. J. Biochem. 196:1-10;  
Coughlin, S.R. (1994) Curr. Opin. Cell Biol. 6:191-197). The amino-terminus of a GPCR is  
extracellular, is of variable length, and is often glycosylated. The carboxy-terminus is cytoplasmic  
25 and generally phosphorylated. Extracellular loops alternate with intracellular loops and link the  
transmembrane domains. Cysteine disulfide bridges linking the second and third extracellular loops  
may interact with agonists and antagonists. The most conserved domains of GPCRs are the  
transmembrane domains and the first two cytoplasmic loops. The transmembrane domains account,  
in part, for structural and functional features of the receptor. In most cases, the bundle of  $\alpha$  helices  
30 forms a ligand-binding pocket. The extracellular N-terminal segment, or one or more of the three  
extracellular loops, may also participate in ligand binding. Ligand binding activates the receptor by  
inducing a conformational change in intracellular portions of the receptor. In turn, the large, third  
intracellular loop of the activated receptor interacts with a heterotrimeric guanine nucleotide binding  
(G) protein complex which mediates further intracellular signaling activities, including the activation  
35 of second messengers such as cyclic AMP (cAMP), phospholipase C, and inositol triphosphate, and

the interaction of the activated GPCR with ion channel proteins. (See, e.g., Watson, S. and S. Arkinstall (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 2-6; Bolander, F.F. (1994) Molecular Endocrinology, Academic Press, San Diego CA, pp. 162-176; Baldwin, J.M. (1994) *Curr. Opin. Cell Biol.* 6:180-190.)

5 GPCRs include receptors for sensory signal mediators (e.g., light and olfactory stimulatory molecules); adenosine,  $\gamma$ -aminobutyric acid (GABA), hepatocyte growth factor, melanocortins, neuropeptide Y, opioid peptides, opsins, somatostatin, tachykinins, vasoactive intestinal polypeptide family, and vasopressin; biogenic amines (e.g., dopamine, epinephrine and norepinephrine, histamine, glutamate (metabotropic effect), acetylcholine (muscarinic effect), and serotonin); chemokines; lipid  
10 mediators of inflammation (e.g., prostaglandins and prostanoids, platelet activating factor, and leukotrienes); and peptide hormones (e.g., bombesin, bradykinin, calcitonin, C5a anaphylatoxin, endothelin, follicle-stimulating hormone (FSH), gonadotropic-releasing hormone (GnRH), neurokinin, and thyrotropin-releasing hormone (TRH), and oxytocin). GPCRs which act as receptors for stimuli that have yet to be identified are known as orphan receptors.

15 The diversity of the GPCR family is further increased by alternative splicing. Many GPCR genes contain introns, and there are currently over 30 such receptors for which splice variants have been identified. The largest number of variations are at the protein C-terminus. N-terminal and cytoplasmic loop variants are also frequent, while variants in the extracellular loops or transmembrane domains are less common. Some receptors have more than one site at which variance  
20 can occur. The splicing variants appear to be functionally distinct, based upon observed differences in distribution, signaling, coupling, regulation, and ligand binding profiles (Kilpatrick, G.J. et al. (1999) *Trends Pharmacol. Sci.* 20:294-301).

GPCRs can be divided into three major subfamilies: the rhodopsin-like, secretin-like, and metabotropic glutamate receptor subfamilies. Members of these GPCR subfamilies share similar  
25 functions and the characteristic seven transmembrane structure, but have divergent amino acid sequences. The largest family consists of the rhodopsin-like GPCRs, which transmit diverse extracellular signals including hormones, neurotransmitters, and light. Rhodopsin is a photosensitive GPCR found in animal retinas. In vertebrates, rhodopsin molecules are embedded in membranous stacks found in photoreceptor (rod) cells. Each rhodopsin molecule responds to a photon of light by  
30 triggering a decrease in cGMP levels which leads to the closure of plasma membrane sodium channels. In this manner, a visual signal is converted to a neural impulse. Other rhodopsin-like GPCRs are directly involved in responding to neurotransmitters. These GPCRs include the receptors for adrenaline (adrenergic receptors), acetylcholine (muscarinic receptors), adenosine, galanin, and glutamate (N-methyl-D-aspartate/NMDA receptors). (Reviewed in Watson, S. and S. Arkinstall  
35 (1994) The G-Protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 7-9, 19-22,

32-35, 130-131, 214-216, 221-222; Habert-Ortoli, E. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9780-9783.)

The galanin receptors mediate the activity of the neuroendocrine peptide galanin, which inhibits secretion of insulin, acetylcholine, serotonin and noradrenaline, and stimulates prolactin and growth hormone release. Galanin receptors are involved in feeding disorders, pain, depression, and Alzheimer's disease (Kask, K. et al. (1997) Life Sci. 60:1523-1533). Other nervous system rhodopsin-like GPCRs include a growing family of receptors for lysophosphatidic acid and other lysophospholipids, which appear to have roles in development and neuropathology (Chun, J. et al. (1999) Cell Biochem. Biophys. 30:213-242).

The largest subfamily of GPCRs, the olfactory receptors, are also members of the rhodopsin-like GPCR family. These receptors function by transducing odorant signals. Numerous distinct olfactory receptors are required to distinguish different odors. Each olfactory sensory neuron expresses only one type of olfactory receptor, and distinct spatial zones of neurons expressing distinct receptors are found in nasal passages. For example, the RA1c receptor which was isolated from a rat brain library, has been shown to be limited in expression to very distinct regions of the brain and a defined zone of the olfactory epithelium (Raming, K. et al. (1998) Receptors Channels 6:141-151). However, the expression of olfactory-like receptors is not confined to olfactory tissues. For example, three rat genes encoding olfactory-like receptors having typical GPCR characteristics showed expression patterns not only in taste and olfactory tissue, but also in male reproductive tissue (Thomas, M.B. et al. (1996) Gene 178:1-5).

Members of the secretin-like GPCR subfamily have as their ligands peptide hormones such as secretin, calcitonin, glucagon, growth hormone-releasing hormone, parathyroid hormone, and vasoactive intestinal peptide. For example, the secretin receptor responds to secretin, a peptide hormone that stimulates the secretion of enzymes and ions in the pancreas and small intestine (Watson, *supra*, pp. 278-283). Secretin receptors are about 450 amino acids in length and are found in the plasma membrane of gastrointestinal cells. Binding of secretin to its receptor stimulates the production of cAMP.

Examples of secretin-like GPCRs implicated in inflammation and the immune response include the EGF module-containing, mucin-like hormone receptor (Emr1) and CD97 receptor proteins. These GPCRs are members of the recently characterized EGF-TM7 receptors subfamily. These seven transmembrane hormone receptors exist as heterodimers *in vivo* and contain between three and seven potential calcium-binding EGF-like motifs. CD97 is predominantly expressed in leukocytes and is markedly upregulated on activated B and T cells (McKnight, A.J. and S. Gordon (1998) J. Leukoc. Biol. 63:271-280).

The third GPCR subfamily is the metabotropic glutamate receptor family. Glutamate is the



major excitatory neurotransmitter in the central nervous system. The metabotropic glutamate receptors modulate the activity of intracellular effectors, and are involved in long-term potentiation (Watson, supra, p.130). The  $\text{Ca}^{2+}$ -sensing receptor, which senses changes in the extracellular concentration of calcium ions, has a large extracellular domain including clusters of acidic amino acids which may be involved in calcium binding. The metabotropic glutamate receptor family also includes pheromone receptors, the  $\text{GABA}_B$  receptors, and the taste receptors.

Other subfamilies of GPCRs include two groups of chemoreceptor genes found in the nematodes Caenorhabditis elegans and Caenorhabditis briggsae, which are distantly related to the mammalian olfactory receptor genes. The yeast pheromone receptors STE2 and STE3, involved in the response to mating factors on the cell membrane, have their own seven-transmembrane signature, as do the cAMP receptors from the slime mold Dictyostelium discoideum, which are thought to regulate the aggregation of individual cells and control the expression of numerous developmentally-regulated genes.

GPCR mutations, which may cause loss of function or constitutive activation, have been associated with numerous human diseases (Coughlin, supra). For instance, retinitis pigmentosa may arise from mutations in the rhodopsin gene. Furthermore, somatic activating mutations in the thyrotropin receptor have been reported to cause hyperfunctioning thyroid adenomas, suggesting that certain GPCRs susceptible to constitutive activation may behave as protooncogenes (Parma, J. et al. (1993) *Nature* 365:649-651). GPCR receptors for the following ligands also contain mutations associated with human disease: luteinizing hormone (precocious puberty); vasopressin  $\text{V}_2$  (X-linked nephrogenic diabetes); glucagon (diabetes and hypertension); calcium (hyperparathyroidism, hypocalcuria, hypercalcemia); parathyroid hormone (short limbed dwarfism);  $\beta_3$ -adrenoceptor (obesity, non-insulin-dependent diabetes mellitus); growth hormone releasing hormone (dwarfism); and adrenocorticotropin (glucocorticoid deficiency) (Wilson, S. et al. (1998) *Br. J. Pharmacol.* 125:1387-1392; Stadel, J.M. et al. (1997) *Trends Pharmacol. Sci.* 18:430-437). GPCRs are also involved in depression, schizophrenia, sleeplessness, hypertension, anxiety, stress, renal failure, and several cardiovascular disorders (Horn, F. and G. Vriend (1998) *J. Mol. Med.* 76:464-468).

In addition, within the past 20 years several hundred new drugs have been recognized that are directed towards activating or inhibiting GPCRs. The therapeutic targets of these drugs span a wide range of diseases and disorders, including cardiovascular, gastrointestinal, and central nervous system disorders as well as cancer, osteoporosis and endometriosis (Wilson, supra; Stadel, supra). For example, the dopamine agonist L-dopa is used to treat Parkinson's disease, while a dopamine antagonist is used to treat schizophrenia and the early stages of Huntington's disease. Agonists and antagonists of adrenoceptors have been used for the treatment of asthma, high blood pressure, other cardiovascular disorders, and anxiety; muscarinic agonists are used in the treatment of glaucoma and

tachycardia; serotonin 5HT<sub>1D</sub> antagonists are used against migraine; and histamine H<sub>1</sub> antagonists are used against allergic and anaphylactic reactions, hay fever, itching, and motion sickness (Horn, supra).

Recent research suggests potential future therapeutic uses for GPCRs in the treatment of metabolic disorders including diabetes, obesity, and osteoporosis. For example, mutant V2 vasopressin receptors causing nephrogenic diabetes could be functionally rescued in vitro by co-expression of a C-terminal V2 receptor peptide spanning the region containing the mutations. This result suggests a possible novel strategy for disease treatment (Schöneberg, T. et al. (1996) EMBO J. 15:1283-1291). Mutations in melanocortin-4 receptor (MC4R) are implicated in human weight regulation and obesity. As with the vasopressin V2 receptor mutants, these MC4R mutants are defective in trafficking to the plasma membrane (Ho, G. and R.G. MacKenzie (1999) J. Biol. Chem. 274:35816-35822), and thus might be treated with a similar strategy. The type 1 receptor for parathyroid hormone (PTH) is a GPCR that mediates the PTH-dependent regulation of calcium homeostasis in the bloodstream. Study of PTH/receptor interactions may enable the development of novel PTH receptor ligands for the treatment of osteoporosis (Mannstadt, M. et al. (1999) Am. J. Physiol. 277:F665-F675).

The chemokine receptor group of GPCRs have potential therapeutic utility in inflammation and infectious disease. (For review, see Locati, M. and P.M. Murphy (1999) Annu. Rev. Med. 50:425-440.) Chemokines are small polypeptides that act as intracellular signals in the regulation of leukocyte trafficking, hematopoiesis, and angiogenesis. Targeted disruption of various chemokine receptors in mice indicates that these receptors play roles in pathologic inflammation and in autoimmune disorders such as multiple sclerosis. Chemokine receptors are also exploited by infectious agents, including herpesviruses and the human immunodeficiency virus (HIV-1) to facilitate infection. A truncated version of chemokine receptor CCR5, which acts as a coreceptor for infection of T-cells by HIV-1, results in resistance to AIDS, suggesting that CCR5 antagonists could be useful in preventing the development of AIDS.

The discovery of new G-protein coupled receptors and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of G-protein coupled receptors.

## SUMMARY OF THE INVENTION

The invention features purified polypeptides, G-protein coupled receptors, referred to

collectively as "GCREC" and individually as "GCREC-1," "GCREC-2," "GCREC-3," "GCREC-4," "GCREC-5," "GCREC-6," "GCREC-7," and "GCREC-8." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-8.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-8. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:9-16.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group

consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting

of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide  
5 complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a  
10 polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic  
15 fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment the  
20 composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence  
25 selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the  
30 invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as  
35 an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an

amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional GCREC, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in

altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:9-16, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

5           The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, ii) a  
10 naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological  
15 sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, ii) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide  
20 of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated  
25 biological sample is indicative of toxicity of the test compound.

### BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

30           Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for  
35 analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

5 Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

10 Table 8 shows tissue-specific expression of polynucleotides of the invention.

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these  
15 may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a  
20 reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.  
25 Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the  
30 invention is not entitled to antedate such disclosure by virtue of prior invention.

### DEFINITIONS

"GCREC" refers to the amino acid sequences of substantially purified GCREC obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

35 The term "agonist" refers to a molecule which intensifies or mimics the biological activity of



GCREC. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GCREC either by directly interacting with GCREC or by acting on components of the biological pathway in which GCREC participates.

An "allelic variant" is an alternative form of the gene encoding GCREC. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding GCREC include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as GCREC or a polypeptide with at least one functional characteristic of GCREC. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GCREC, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GCREC. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GCREC. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GCREC is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity

of GCREC. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GCREC either by directly interacting with GCREC or by acting on components of the biological pathway in which GCREC participates.

5           The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind GCREC polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the  
10 translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and Keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

          The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that  
15 makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

20           The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having  
25 modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the  
30 designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

          The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic GCREC, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific  
35 antibodies.

“Complementary” describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A “composition comprising a given polynucleotide sequence” and a “composition comprising a given amino acid sequence” refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GCREC or fragments of GCREC may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

“Consensus sequence” refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

“Conservative amino acid substitutions” are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
25	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
30	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
35	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
40	Thr	Ser, Val

Trp  
Tyr  
Val

Phe, Tyr  
His, Phe, Trp  
Ile, Leu, Thr

---

5           Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

10           A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

          The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative  
15   polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

          A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

20           "Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

          A "fragment" is a unique portion of GCREC or the polynucleotide encoding GCREC which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up  
25   to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For  
30   example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

35           A fragment of SEQ ID NO:9-16 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:9-16, for example, as distinct from any other sequence in the

genome from which the fragment was obtained. A fragment of SEQ ID NO:9-16 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:9-16 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:9-16 and the region of SEQ ID NO:9-16 to which the fragment corresponds are routinely  
5 determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-8 is encoded by a fragment of SEQ ID NO:9-16. A fragment of SEQ ID NO:1-8 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-8. For example, a fragment of SEQ ID NO:1-8 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-8. The precise length of a  
10 fragment of SEQ ID NO:1-8 and the region of SEQ ID NO:1-8 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A  
15 "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a  
20 standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e  
25 sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue  
30 weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available  
35 from several sources, including the NCBI, Bethesda, MD, and on the Internet at

http://www.ncbi.nlm.nih.gov/BLAST/. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

10        *Matrix: BLOSUM62*  
           *Reward for match: 1*  
           *Penalty for mismatch: -2*  
           *Open Gap: 5 and Extension Gap: 2 penalties*  
           *Gap x drop-off: 50*  
 15        *Expect: 10*  
           *Word Size: 11*  
           *Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

25        Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

30        The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

35        Percent identity between polypeptide sequences may be determined using the default

parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default  
5 residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for  
10 example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

15 *Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for  
20 instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain  
25 DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

30 "Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the  
35 stringency of the hybridization process, with more stringent conditions allowing less non-specific

binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity.

- 5 Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

- 15 High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

25 The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

- 35 "Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression



of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of GCREC which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of GCREC which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of GCREC. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GCREC.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an GCREC may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of GCREC.

"Probe" refers to nucleic acid sequences encoding GCREC, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule.

Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

“Primers” are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and  
5 identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers  
10 may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular  
15 Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the  
20 PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a “mispriming library,” in which  
25 sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user’s specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments,  
30 thereby allowing selection of primers that hybridize to either the most conserved or least conserved

regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to  
5 identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the  
10 artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to  
15 transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated  
20 regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent,  
25 chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose  
30 instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing GCREC, nucleic acids encoding GCREC, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

35 The terms "specific binding" and "specifically binding" refer to that interaction between a

protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide  
5 comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which  
10 they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers,  
15 microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient  
20 cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term  
25 "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic  
30 acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The  
35 transgenic organisms contemplated in accordance with the present invention include bacteria,

cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989),

5 supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

## THE INVENTION

The invention is based on the discovery of new human G-protein coupled receptors (GCREC), the polynucleotides encoding GCREC, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, neurological, cardiovascular, gastrointestinal,

autoimmune/inflammatory, and metabolic disorders, and viral infections.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are G-protein coupled receptors. For example, SEQ ID NO:2 is 36% identical to mouse P2Y1 receptor (GenBank ID g6013075) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.0e-55, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:2 also contains a seven transmembrane receptor (rhodopsin family) domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:2 is a G-protein coupled receptor. In an alternative example, SEQ ID NO:3 is 52% identical to HM74

(GenBank ID g219867) as determined by BLAST. (See Table 2.) The BLAST probability score is 3.7e-88. Data from BLAST-DOMO, MOTIFS, BLIMPS-BLOCKS, BLIMPS-PRINTS, PROFILESCAN, HMMER-PFAM, and SPSCAN analyses provide further corroborative evidence that SEQ ID NO:3 is a G-protein coupled receptor. In an alternative example, SEQ ID NO:8 is 50% identical to a predicted mouse odorant receptor-like protein (GenBank ID g6532001) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 8.4e-75, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:8 also contains a seven transmembrane receptor (rhodopsin family) domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:8 is a G-protein coupled receptor. SEQ ID NO:1 and SEQ ID NO:4-7 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-8 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:9-16 or that distinguish between SEQ ID NO:9-16 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 6251915H1 is the identification number of an Incyte cDNA sequence, and LUNPTUT02 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 70812700V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5663306) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to

coding regions predicted by Genscan analysis of genomic DNA. For example, GNN.g7528005\_000007\_004 is the identification number of a Genscan-predicted coding sequence, with g7528005 being the GenBank identification number of the sequence to which Genscan was applied. The Genscan-predicted coding sequences may have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon-stretching" algorithm. (See Example V.) In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

Table 8 shows tissue-specific expression of polynucleotides of the invention. Column 1 lists groups of tissues which were tested by polymerase chain reaction (PCR) for expression of the polynucleotides. The remaining columns indicate whether a particular polynucleotide was expressed in each tissue group. Detection of a PCR product indicated positive expression, denoted by a "+" sign, while inability to detect a PCR product indicated a lack of expression, denoted by a "-" sign.

The invention also encompasses GCREC variants. A preferred GCREC variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the GCREC amino acid sequence, and which contains at least one functional or structural characteristic of GCREC.

The invention also encompasses polynucleotides which encode GCREC. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:9-16, which encodes GCREC. The polynucleotide sequences of SEQ ID NO:9-16, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding GCREC. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GCREC. A particular aspect of the invention encompasses a variant of a



polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:9-16 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:9-16. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GCREC.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GCREC, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GCREC, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GCREC and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring GCREC under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GCREC or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GCREC and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GCREC and GCREC derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GCREC or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:9-16 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of

the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE  
5 amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA  
10 sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GCREC may be extended utilizing a partial nucleotide  
15 sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown  
20 sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme  
25 digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries  
30 and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

35 When screening for full length cDNAs, it is preferable to use libraries that have been

size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

5           Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate  
10       software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

          In another embodiment of the invention, polynucleotide sequences or fragments thereof  
15       which encode GCREC may be cloned in recombinant DNA molecules that direct expression of GCREC, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GCREC.

          The nucleotide sequences of the present invention can be engineered using methods generally  
20       known in the art in order to alter GCREC-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction  
25       sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

          The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or  
30       improve the biological properties of GCREC, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of  
35       DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial"

breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby  
5 maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding GCREC may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.)  
10 Alternatively, GCREC itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the  
15 amino acid sequence of GCREC, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.)  
20 The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active GCREC, the nucleotide sequences encoding GCREC or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding  
25 sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GCREC. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GCREC. Such signals include the ATG initiation codon and adjacent sequences,  
30 e.g. the Kozak sequence. In cases where sequences encoding GCREC and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and  
35 initiation codons may be of various origins, both natural and synthetic. The efficiency of expression

may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GCRC and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GCRC. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GCRC. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GCRC can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSORT1 plasmid (Life Technologies). Ligation of sequences encoding GCRC into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of

nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GCREC are needed, e.g. for the production of antibodies, vectors which direct high level expression of GCREC may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

5 Yeast expression systems may be used for production of GCREC. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel,  
10 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GCREC. Transcription of sequences encoding GCREC may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.  
15 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology  
20 (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GCREC may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain  
25 infective virus which expresses GCREC in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of  
30 DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression  
35 of GCREC in cell lines is preferred. For example, sequences encoding GCREC can be transformed

into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap<sup>r</sup>* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GCREC is inserted within a marker gene sequence, transformed cells containing sequences encoding GCREC can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GCREC under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GCREC and that express GCREC may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GCREC using either

specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GCREC is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GCREC include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GCREC, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GCREC may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GCREC may be designed to contain signal sequences which direct secretion of GCREC through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct



modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GCREC may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GCREC protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GCREC activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GCREC encoding sequence and the heterologous protein sequence, so that GCREC may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GCREC may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

GCREC of the present invention or fragments thereof may be used to screen for compounds that specifically bind to GCREC. At least one and up to a plurality of test compounds may be screened for specific binding to GCREC. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of GCREC, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which GCREC binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express GCREC, either as a secreted

protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing GCREC or cell membrane fractions which contain GCREC are then contacted with a test compound and binding, stimulation, or inhibition of activity of either GCREC or the compound is analyzed.

5           An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with GCREC, either in solution or affixed to a solid support, and detecting the binding of GCREC to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a  
10   labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

          GCREC of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of GCREC. Such compounds may include agonists, antagonists, or partial  
15   or inverse agonists. In one embodiment, an assay is performed under conditions permissive for GCREC activity, wherein GCREC is combined with at least one test compound, and the activity of GCREC in the presence of a test compound is compared with the activity of GCREC in the absence of the test compound. A change in the activity of GCREC in the presence of the test compound is indicative of a compound that modulates the activity of GCREC. Alternatively, a test compound is  
20   combined with an in vitro or cell-free system comprising GCREC under conditions suitable for GCREC activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of GCREC may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

          In another embodiment, polynucleotides encoding GCREC or their mammalian homologs  
25   may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the  
30   gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic  
35   Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell

blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

5 Polynucleotides encoding GCREC may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

10 Polynucleotides encoding GCREC can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding GCREC is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and  
15 treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress GCREC, e.g., by secreting GCREC in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

## THERAPEUTICS

20 Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GCREC and G-protein coupled receptors. In addition, the expression of GCREC is closely associated with brain tumor, breast tumor, liver, fetal kidney, and fetal thymus tissue. Therefore, GCREC appears to play a role in cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections. In the  
25 treatment of disorders associated with increased GCREC expression or activity, it is desirable to decrease the expression or activity of GCREC. In the treatment of disorders associated with decreased GCREC expression or activity, it is desirable to increase the expression or activity of GCREC.

Therefore, in one embodiment, GCREC or a fragment or derivative thereof may be  
30 administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma,  
35 leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of

the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic

carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired

5 immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and

10 hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis,

15 cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis,

20 polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a metabolic disorder such as diabetes, obesity, and osteoporosis; and an infection by a viral

25 agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, and tongavirus.

In another embodiment, a vector capable of expressing GCREC or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased

30 expression or activity of GCREC including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified GCREC in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GCREC may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC including, but not limited to, those listed above.

In a further embodiment, an antagonist of GCREC may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCREC. Examples of such disorders include, but are not limited to, those cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections, described above. In one aspect, an antibody which specifically binds GCREC may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express GCREC.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GCREC may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCREC including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GCREC may be produced using methods which are generally known in the art. In particular, purified GCREC may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GCREC. Antibodies to GCREC may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with GCREC or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to

GCREC have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of GCREC amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GCREC may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce GCREC-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for GCREC may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GCREC and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies

reactive to two non-interfering GCREC epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GCREC. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of GCREC-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GCREC epitopes, represents the average affinity, or avidity, of the antibodies for GCREC. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular GCREC epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the GCREC-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GCREC, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of GCREC-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding GCREC, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding GCREC. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GCREC. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence



complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

10 In another embodiment of the invention, polynucleotides encoding GCREC may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency 15 (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) 20 express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides 25 brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in GCREC expression or regulation causes disease, the expression of GCREC from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in 30 GCREC are treated by constructing mammalian expression vectors encoding GCREC and introducing these vectors by mechanical means into GCREC-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev.* 35 *Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr.*

Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of GCREC include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). GCREC may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) Proc. Natl. Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. supra)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding GCREC from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) Virology 52:456-467), or by electroporation (Neumann, E. et al. (1982) EMBO J. 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to GCREC expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding GCREC under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant")

discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding GCREC to cells which have one or more genetic abnormalities with respect to the expression of GCREC. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding GCREC to target cells which have one or more genetic abnormalities with respect to the expression of GCREC. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing GCREC to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of

ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding GCREC to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based  
5 on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for GCREC into the  
10 alphavirus genome in place of the capsid-coding region results in the production of a large number of GCREC-coding RNAs and the synthesis of high levels of GCREC in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy  
15 application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of GCREC into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the  
20 art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of  
25 polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

30 Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GCREC.

35 Specific ribozyme cleavage sites within any potential RNA target are initially identified by

scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GCREC. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding GCREC. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased GCREC expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding GCREC may be therapeutically useful, and in the treatment of disorders associated with decreased GCREC expression or activity, a compound which specifically promotes expression of the polynucleotide encoding GCREC may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in

altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound  
5 based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding GCREC is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding GCREC are assayed  
10 by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding GCREC. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide  
15 exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

25 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.  
30 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition  
35 which generally comprises an active ingredient formulated with a pharmaceutically acceptable

excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of GCREC, antibodies to GCREC, and mimetics, agonists, antagonists, or inhibitors of

5 GCREC.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

10 Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the

15 lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination

20 of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising GCREC or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, GCREC or a fragment thereof may be joined to a short cationic N-

25 terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs,

30 monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GCREC or fragments thereof, antibodies of GCREC, and agonists, antagonists or inhibitors of

35 GCREC, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be

determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large  
5 therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

10 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and  
15 response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

20 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind GCREC may be used for the  
25 diagnosis of disorders characterized by expression of GCREC, or in assays to monitor patients being treated with GCREC or agonists, antagonists, or inhibitors of GCREC. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GCREC include methods which utilize the antibody and a label to detect GCREC in human body fluids or in extracts of cells or tissues. The antibodies may be used with or  
30 without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GCREC, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GCREC expression.

35 Normal or standard values for GCREC expression are established by combining body fluids or cell



extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to GCREC under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of GCREC expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for  
5 diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GCREC may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect  
10 and quantify gene expression in biopsied tissues in which expression of GCREC may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of GCREC, and to monitor regulation of GCREC levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GCREC or closely related molecules may be used  
15 to identify nucleic acid sequences which encode GCREC. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding GCREC, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the GCREC encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:9-16 or from genomic sequences including promoters, enhancers, and introns of the GCREC gene.  
20

Means for producing specific hybridization probes for DNAs encoding GCREC include the cloning of polynucleotide sequences encoding GCREC or GCREC derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels,  
25 such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.  
30

Polynucleotide sequences encoding GCREC may be used for the diagnosis of disorders associated with expression of GCREC. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal  
35 hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including

adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a

5 neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural

10 abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central

15 nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia,

20 catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, complications of thrombolysis, balloon

25 angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus

30 erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal

obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic

5 obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-

10 occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis,

15 autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis,

20 myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and

25 trauma; a metabolic disorder such as diabetes, obesity, and osteoporosis; and an infection by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, and tongavirus. The polynucleotide sequences encoding GCREC may be used in Southern or northern analysis, dot blot, or other

30 membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered GCREC expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GCREC may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide

35 sequences encoding GCREC may be labeled by standard methods and added to a fluid or tissue

sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GCREC in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GCREC, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GCREC, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GCREC may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding GCREC, or a fragment of a polynucleotide complementary to the polynucleotide encoding GCREC, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences

encoding GCREC may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, 5 oligonucleotide primers derived from the polynucleotide sequences encoding GCREC are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the 10 oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed in silico SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation 15 of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of GCREC include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from 20 standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

25 In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene 30 function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly 35 effective and display the fewest side effects may be selected for a patient based on his/her

pharmacogenomic profile.

In another embodiment, GCREC, fragments of GCREC, or antibodies specific for GCREC may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described  
5 above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at  
10 a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present  
15 invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

20 Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and  
25 toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide  
30 measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms,  
35 knowledge of gene function is not necessary for the statistical matching of signatures which leads to

prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

5           In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with  
10   levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

          Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a  
15   proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are  
20   separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The  
25   optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by  
30   comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

          A proteomic profile may also be generated using antibodies specific for GCREC to quantify the levels of GCREC expression. In one embodiment, the antibodies are used as elements on a  
35   microarray, and protein expression levels are quantified by exposing the microarray to the sample and

detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) Anal. Biochem. 270:103-111; Mendozze, L.G. et al. (1999) Biotechniques 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) Electrophoresis 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.



In another embodiment of the invention, nucleic acid sequences encoding GCREC may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding GCREC on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, GCREC, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GCREC and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with GCREC, or fragments thereof, and washed. Bound GCREC is then detected by methods well known in the art. Purified GCREC can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GCREC specifically compete with a test compound for binding GCREC. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GCREC.

In additional embodiments, the nucleotide sequences which encode GCREC may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, including U.S. Ser. No. 60/205,628, U.S. Ser. No. 60/207,556, U.S. Ser. No. 60/208,834, U.S. Ser. No. 60/206,222, and U.S. Ser. No. 60/208,861, are expressly incorporated by reference herein.

## EXAMPLES

### I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal  
5 cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides  
10 were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques  
15 disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public  
20 databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The  
25 Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on  
30 GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite,  
35 and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length

polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:9-16. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

#### **IV. Identification and Editing of Coding Sequences from Genomic DNA**

Putative G-protein coupled receptors were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpr and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode G-protein coupled receptors, the encoded polypeptides were analyzed by querying against PFAM models for G-protein coupled receptors. Potential G-protein coupled receptors were also identified by homology to Incyte cDNA sequences that had been annotated as G-protein coupled receptors. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpr public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available,

this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

#### **V. Assembly of Genomic Sequence Data with cDNA Sequence Data**

##### **"Stitched" Sequences**

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

##### **"Stretched" Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The

GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

## 5 VI. Chromosomal Mapping of GCREC Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:9-16 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:9-16 were assembled into clusters of contiguous and overlapping sequences using  
10 assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Génethon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

15 Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM  
20 distances are based on genetic markers mapped by Génethon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

## 25 VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel (1995) supra, ch. 4 and 16.)

30 Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding GCREC are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding GCREC. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

### **VIII. Extension of GCREC Encoding Polynucleotides**

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was



synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was  
5 quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

10 In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### **IX. Labeling and Use of Individual Hybridization Probes**

Hybridization probes derived from SEQ ID NO:9-16 are employed to screen cDNAs,  
15 genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase  
20 (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

25 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and  
30 compared.

#### **X. Microarrays**

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the  
35 aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999),

supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### **Tissue or Cell Sample Preparation**

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### **Microarray Preparation**

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5  
5  $\mu$ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR  
10 Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1  $\mu$ l of the array element DNA, at an average  
15 concentration of 100 ng/ $\mu$ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate  
20 buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

### **Hybridization**

Hybridization reactions contain 9  $\mu$ l of sample mixture consisting of 0.2  $\mu$ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample  
25 mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140  $\mu$ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash  
30 buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

### **Detection**

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines

at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### **XI. Complementary Polynucleotides**

Sequences complementary to the GCRC-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GCRC. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same

procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GCREC. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GCREC-encoding transcript.

## XII. Expression of GCREC

Expression and purification of GCREC is achieved using bacterial or virus-based expression systems. For expression of GCREC in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GCREC upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GCREC in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GCREC by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, GCREC is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from GCREC at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified GCREC obtained by these methods can be used directly in the assays

shown in Examples XVI, XVII, and XVIII, where applicable.

### **XIII. Functional Assays**

GCREC function is assessed by expressing the sequences encoding GCREC at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GCREC on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GCREC and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GCREC and other genes of interest can be analyzed by northern analysis or microarray techniques.

### **XIV. Production of GCREC Specific Antibodies**

GCREC substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the GCREC amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-GCREC activity by, for example, binding the peptide or GCREC to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

#### **XV. Purification of Naturally Occurring GCREC Using Specific Antibodies**

Naturally occurring or recombinant GCREC is substantially purified by immunoaffinity chromatography using antibodies specific for GCREC. An immunoaffinity column is constructed by covalently coupling anti-GCREC antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GCREC are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GCREC (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GCREC binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GCREC is collected.

#### **XVI. Identification of Molecules Which Interact with GCREC**

Molecules which interact with GCREC may include agonists and antagonists, as well as molecules involved in signal transduction, such as G proteins. GCREC, or a fragment thereof, is labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) A fragment of GCREC includes, for example, a fragment comprising one or more of the three extracellular loops, the extracellular N-terminal region, or the third intracellular loop. Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GCREC, washed, and any wells with labeled GCREC complex are assayed. Data obtained using different concentrations of GCREC are used to calculate values for the number, affinity, and association of GCREC with the candidate ligand molecules.

Alternatively, molecules interacting with GCREC are analyzed using the yeast two-hybrid



system as described in Fields, S. and O. Song (1989) Nature 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech). GCREC may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions  
5 between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Potential GCREC agonists or antagonists may be tested for activation or inhibition of GCREC receptor activity using the assays described in sections XVII and XVIII. Candidate molecules may be selected from known GPCR agonists or antagonists, peptide libraries, or  
10 combinatorial chemical libraries.

Methods for detecting interactions of GCREC with intracellular signal transduction molecules such as G proteins are based on the premise that internal segments or cytoplasmic domains from an orphan G protein-coupled seven transmembrane receptor may be exchanged with the analogous domains of a known G protein-coupled seven transmembrane receptor and used to identify  
15 the G-proteins and downstream signaling pathways activated by the orphan receptor domains (Kobilka, B.K. et al. (1988) Science 240:1310-1316). In an analogous fashion, domains of the orphan receptor may be cloned as a portion of a fusion protein and used in binding assays to demonstrate interactions with specific G proteins. Studies have shown that the third intracellular loop of G protein-coupled seven transmembrane receptors is important for G protein interaction and signal  
20 transduction (Conklin, B.R. et al. (1993) Cell 73:631-641). For example, the DNA fragment corresponding to the third intracellular loop of GCREC may be amplified by the polymerase chain reaction (PCR) and subcloned into a fusion vector such as pGEX (Pharmacia Biotech). The construct is transformed into an appropriate bacterial host, induced, and the fusion protein is purified from the cell lysate by glutathione-Sepharose 4B (Pharmacia Biotech) affinity chromatography.

25 For in vitro binding assays, cell extracts containing G proteins are prepared by extraction with 50 mM Tris, pH 7.8, 1 mM EGTA, 5 mM MgCl<sub>2</sub>, 20 mM CHAPS, 20% glycerol, 10 µg of both aprotinin and leupeptin, and 20 µl of 50 mM phenylmethylsulfonyl fluoride. The lysate is incubated on ice for 45 min with constant stirring, centrifuged at 23,000 g for 15 min at 4°C, and the supernatant is collected. 750 µg of cell extract is incubated with glutathione S-transferase (GST)  
30 fusion protein beads for 2 h at 4°C. The GST beads are washed five times with phosphate-buffered saline. Bound G subunits are detected by [<sup>32</sup>P]ADP-ribosylation with pertussis or cholera toxins. The reactions are terminated by the addition of SDS sample buffer (4.6% (w/v) SDS, 10% (v/v) β-mercaptoethanol, 20% (w/v) glycerol, 95.2 mM Tris-HCl, pH 6.8, 0.01% (w/v) bromphenol blue). The [<sup>32</sup>P]ADP-labeled proteins are separated on 10% SDS-PAGE gels, and autoradiographed. The  
35 separated proteins in these gels are transferred to nitrocellulose paper, blocked with blotto (5% nonfat

dried milk, 50 mM Tris-HCl (pH 8.0), 2 mM CaCl<sub>2</sub>, 80 mM NaCl, 0.02% NaN<sub>3</sub>, and 0.2% Nonidet P-40) for 1 hour at room temperature, followed by incubation for 1.5 hours with G $\alpha$  subtype selective antibodies (1:500; Calbiochem-Novabiochem). After three washes, blots are incubated with horseradish peroxidase (HRP)-conjugated goat anti-rabbit immunoglobulin (1:2000, Cappel, Westchester PA) and visualized by the chemiluminescence-based ECL method (Amersham Corp.).

#### XVII. Demonstration of GCREC Activity

An assay for GCREC activity measures the expression of GCREC on the cell surface. cDNA encoding GCREC is transfected into an appropriate mammalian cell line. Cell surface proteins are labeled with biotin as described (de la Fuente, M.A. et al. (1997) Blood 90:2398-2405).

Immunoprecipitations are performed using GCREC-specific antibodies, and immunoprecipitated samples are analyzed using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of GCREC expressed on the cell surface.

In the alternative, an assay for GCREC activity is based on a prototypical assay for ligand/receptor-mediated modulation of cell proliferation. This assay measures the rate of DNA synthesis in Swiss mouse 3T3 cells. A plasmid containing polynucleotides encoding GCREC is added to quiescent 3T3 cultured cells using transfection methods well known in the art. The transiently transfected cells are then incubated in the presence of [<sup>3</sup>H]thymidine, a radioactive DNA precursor molecule. Varying amounts of GCREC ligand are then added to the cultured cells. Incorporation of [<sup>3</sup>H]thymidine into acid-precipitable DNA is measured over an appropriate time interval using a radioisotope counter, and the amount incorporated is directly proportional to the amount of newly synthesized DNA. A linear dose-response curve over at least a hundred-fold GCREC ligand concentration range is indicative of receptor activity. One unit of activity per milliliter is defined as the concentration of GCREC producing a 50% response level, where 100% represents maximal incorporation of [<sup>3</sup>H]thymidine into acid-precipitable DNA (McKay, I. and I. Leigh, eds. (1993) Growth Factors: A Practical Approach, Oxford University Press, New York NY, p. 73.)

In a further alternative, the assay for GCREC activity is based upon the ability of GPCR family proteins to modulate G protein-activated second messenger signal transduction pathways (e.g., cAMP; Gaudin, P. et al. (1998) J. Biol. Chem. 273:4990-4996). A plasmid encoding full length GCREC is transfected into a mammalian cell line (e.g., Chinese hamster ovary (CHO) or human embryonic kidney (HEK-293) cell lines) using methods well-known in the art. Transfected cells are grown in 12-well trays in culture medium for 48 hours, then the culture medium is discarded, and the attached cells are gently washed with PBS. The cells are then incubated in culture medium with or without ligand for 30 minutes, then the medium is removed and cells lysed by treatment with 1 M

perchloric acid. The cAMP levels in the lysate are measured by radioimmunoassay using methods well-known in the art. Changes in the levels of cAMP in the lysate from cells exposed to ligand compared to those without ligand are proportional to the amount of GCREC present in the transfected cells.

5 To measure changes in inositol phosphate levels, the cells are grown in 24-well plates containing  $1 \times 10^5$  cells/well and incubated with inositol-free media and [ $^3\text{H}$ ]myoinositol, 2  $\mu\text{Ci}$ /well, for 48 hr. The culture medium is removed, and the cells washed with buffer containing 10 mM LiCl followed by addition of ligand. The reaction is stopped by addition of perchloric acid. Inositol phosphates are extracted and separated on Dowex AG1-X8 (Bio-Rad) anion exchange resin, and the  
10 total labeled inositol phosphates counted by liquid scintillation. Changes in the levels of labeled inositol phosphate from cells exposed to ligand compared to those without ligand are proportional to the amount of GCREC present in the transfected cells.

#### XVIII. Identification of GCREC Ligands

GCREC is expressed in a eukaryotic cell line such as CHO (Chinese Hamster Ovary) or HEK  
15 (Human Embryonic Kidney) 293 which have a good history of GPCR expression and which contain a wide range of G-proteins allowing for functional coupling of the expressed GCREC to downstream effectors. The transformed cells are assayed for activation of the expressed receptors in the presence of candidate ligands. Activity is measured by changes in intracellular second messengers, such as cyclic AMP or  $\text{Ca}^{2+}$ . These may be measured directly using standard methods well known in the art,  
20 or by the use of reporter gene assays in which a luminescent protein (e.g. firefly luciferase or green fluorescent protein) is under the transcriptional control of a promoter responsive to the stimulation of protein kinase C by the activated receptor (Milligan, G. et al. (1996) Trends Pharmacol. Sci. 17:235-237). Assay technologies are available for both of these second messenger systems to allow high throughput readout in multi-well plate format, such as the adenylyl cyclase activation FlashPlate  
25 Assay (NEN Life Sciences Products), or fluorescent  $\text{Ca}^{2+}$  indicators such as Fluo-4 AM (Molecular Probes) in combination with the FLIPR fluorimetric plate reading system (Molecular Devices). In cases where the physiologically relevant second messenger pathway is not known, GCREC may be coexpressed with the G-proteins  $G_{\alpha 15/16}$  which have been demonstrated to couple to a wide range of G-proteins (Offermanns, S. and M.I. Simon (1995) J. Biol. Chem. 270:15175-15180), in order to funnel  
30 the signal transduction of the GCREC through a pathway involving phospholipase C and  $\text{Ca}^{2+}$  mobilization. Alternatively, GCREC may be expressed in engineered yeast systems which lack endogenous GPCRs, thus providing the advantage of a null background for GCREC activation screening. These yeast systems substitute a human GPCR and  $G_\alpha$  protein for the corresponding components of the endogenous yeast pheromone receptor pathway. Downstream signaling pathways  
35 are also modified so that the normal yeast response to the signal is converted to positive growth on

selective media or to reporter gene expression (Broach, J.R. and J. Thorner (1996) Nature 384 (supp.):14-16). The receptors are screened against putative ligands including known GPCR ligands and other naturally occurring bioactive molecules. Biological extracts from tissues, biological fluids and cell supernatants are also screened.

5           Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention  
10   which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
7474872	1	7474872CD1	9	7474872CB1
6575963	2	6575963CD1	10	6575963CB1
7474846	3	7474846CD1	11	7474846CB1
1258785	4	1258785CD1	12	1258785CB1
1874944	5	1874944CD1	13	1874944CB1
7475270	6	7475270CD1	14	7475270CB1
55000189	7	55000189CD1	15	55000189CB1
7474839	8	7474839CD1	16	7474839CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability Score	GenBank Homolog
1	7474872CD1	g304384	4.10E-53	G protein-coupled receptor [Gallus gallus]
2	6575963CD1	g6013075	1.00E-55	P2Y1 receptor [Mus musculus]
3	7474846CD1	g219867	3.70E-88	HM74 [Homo sapiens] (Nomura, H. et al. (1993) Int. Immunol. 5:1239-1249)
4	1258785CD1	g2894086	2.50E-128	Lanthionine synthetase C-like protein 1 (seven transmembrane-domain protein) [Homo sapiens]
5	1874944CD1	g4164061	8.50E-35	Latrophilin 3 splice variant abbg [Bos taurus] (Matsushita, H. et al. (1999) FEBS Lett. 443:348-352)
6	7475270CD1	g3983394	1.00E-83	Olfactory receptor F7 [Mus musculus] (Krautwurst, D. et al. (1998) Cell 95:917-926)
7	55000189CD1	g1419016	6.90E-106	Odorant receptor [Mus musculus] (Asai, H. et al. (1996) Biochem. Biophys. Res. Commun. 221:240-247)
8	7474839CD1	g11875778	1.00E-100	Prostate specific G-protein coupled receptor; PSGR [Homo sapiens] (Xu, L.L. et al. (2000) Cancer Res. 60:6568-6572)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	7474872CD1	372	T21 S211 S226 T232 T307 S332 S178 T342	N4 N9 N251 N323	Transmembrane domain: V275-L295 Rhodopsin-like GPCR superfamily PR00237: F101-I123; V137-R158; V192-F215; T232-V256; L279-R305; L24-L48; V57-R78 RECEPTOR COUPLED GPROTEIN TRANSMEMBRANE GLYCOPROTEIN PHOSPHORYLATION LIPOPROTEIN PALMITATE PD000009: R52-P155 G-Protein Receptor: G107-I123 G-PROTEIN COUPLED RECEPTORS DM00013 JC4618 24-304: V32-E302; V27-L311 7 transmembrane receptor (rhodopsin family): L39-Y297 G-protein coupled receptor BL00237: W87-P126; F200-S211; Q227-T253; N289-R305 G-protein coupled receptors signature: G98-V145	HMIMER BLIMPS- PRINTS BLAST- PRODOR MOTIFS BLAST-DOMO HMIMER-PFAM
2	6575963CD1	337	S188 S332 T175 T178 T187 T194 T243 T316	N10 N176 N179 N23	G-PROTEIN COUPLED RECEPTORS DM00013 P48042 45-340: K32-R318 G-protein coupled receptor BL00237A: W99-P138, F207-Y218, C234-L260, N294-Q310 Angiotensin II receptor PR00241A: I39-G47 L122-I129 Rhodopsin-like GPCR superfamily signature PR00237C: Y35-I59, S68-L89, F113-I135, C149-F170, N199-I222, A239-I263, Y284-Q310	ProfileScan BLAST-DOMO BLIMPS- BLOCKS BLIMPS- PRINTS BLIMPS- PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
2					G-protein coupled receptors: F111-V157	PROFILES CAN
					transmembrane domain: Y35-I59	HMMER
					7 transmembrane receptor (rhodopsin family) 7tm_1: G50-Y302	HMMER_PFAM
3	7474846CD1	346	S128 S196 S202 S238 S250 S271 S296 S332 T219 T307 T46	N3	G-PROTEIN COUPLED RECEPTORS DM00013 P49019 21-310: D12-L292 G-PROTEIN COUPLED RECEPTORS DM00013 P48042 45-340: M18-F285 G-PROTEIN COUPLED RECEPTORS DM00013 JC4618 24-304: F26-I294 PROBABLE G-PROTEIN-COUPLED RECEPTOR 6H1 FROM T-CELLS DM08033 P32250 1-308: V24-I294 G-protein coupled receptor BL00237: F185-S196, A213-A239, N270-P286, W81-P120 Rhodopsin-like GPCR superfamily PR00237: V17-C41, S50-R71, L95-V117, V131-L152, H177-V200, A218-Y242, L260-P286 G-protein coupled receptors signature: F93-L139 Signal peptide: M1-G32 Transmembrane domain: P20-M44, F93-A110, C137-N154, I222-L244 7 transmembrane receptor (rhodopsin family) 7tm_1: G32-Y278 G-protein coupled receptors signature: G101-V117	BLAST-DOMO BLAST-DOMO BLAST-DOMO BLAST-DOMO BLAST-DOMO BLIMPS-BLOCKS BLIMPS-PRINTS PROFILES CAN SPSCAN HMMER HMMER-PFAM MOTIFS



Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	1258785CD1	432	S149 S216 S221 S398 S428 S429 T118 T128 T200 T227 T266 T296 T37 T418 Y107 Y325 Y371 Y397	N294	PROTEIN TRANSMEMBRANE SEVEN TRANSMEMBRANE DOMAIN P40 PD138974: F45-L426 ATP/GTP-binding site motif A (P-loop): A212-T219 Signal peptide: M1-E27	BLAST- PRODOR  MOTIFS  SPSCAN
5	1874944CD1	240	S196 S219 S223 T124	N211	PROBABLE HORMONE RECEPTOR EMR1 PRECURSOR DM05221 A57172 465-886: L5-S200 G-PROTEIN COUPLED RECEPTORS FAMILY 2 DM00378 P32214 30-474: L4-K186 RECEPTOR TRANSMEMBRANE GPROTEIN COUPLED GLYCOPROTEIN PRECURSOR SIGNAL TYPE POLYPEPTIDE ALTERNATIVE PD000752: L5-K186 G-protein coupled receptor BL00649: S162-T187, M1-L23 Secretin-like GPCR superfamily PR00249: M1-L23, H38-S63, W80-S105, T124-L144, Q154-L175 G-protein coupled receptors family 2 signatures: A81-S200 Transmembrane domain: I79-T101 7 transmembrane receptor (Secretin family) 7tm_2: M1-V179 Signal peptide: M1-A60	BLAST-DOMO  BLAST-DOMO  BLAST- PRODOR  BLIMPS- BLOCKS BLIMPS- PRINTS  PROFILES SCAN  HMMER HMMER-PFAM  SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	7475270CD1	271	S8 S22 S68 S88 S248	N6	Transmembrane domain: L26-L46 7 transmembrane receptor (rhodopsin family) domain: G42-I155; R223-Y247 G-protein coupled receptors signature BL00237: K91-P130; T239-M255 G-protein coupled receptors signature: F104-G147 Olfactory receptor signature PR00245: M60-K81; L195-G210; A231-L242; S248-L262 Rhodopsin-like GPCR superfamily signature PR00237: V27-Y51; M60-K81; F105-V127; K229-M255 OLFACTORY RECEPTOR PROTEIN RECEPTORLIKE GPROTEIN COUPLED TRANSMEMBRANE GLYCOPROTEIN MULTIGENE FAMILY PD149621: T203-R264 G-PROTEIN COUPLED RECEPTORS DM00013 A57069 15-304: F19-F178; W180-K260 G-protein coupled receptors motif: S111-V127	HMMER HMMER-PFAM BLIMPS-BLOCKS PROFILESCAN BLIMPS-PRINTS BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	55000189CD1	276	T192 S68 T136 S137	N5 N190	Transmembrane domains: F31-I49; V205-V224 7 transmembrane receptor (rhodopsin family) domain: G41-P261 G-protein coupled receptors signature BL00237: Q90-P129 Olfactory receptor signature PR00245: M59-Q80; F176-N190; F237-G252 RECEPTOR OLFACTORY PROTEIN RECEPTORLIKE GPROTEIN COUPLED TRANSMEMBRANE GLYCOPROTEIN MULTIGENE FAMILY PD000921: L166-L244 G-PROTEIN COUPLED RECEPTORS DM00013 P30954 29-316: S18-T276 G-PROTEIN COUPLED RECEPTORS: DM00013 P23269 15-304: P118-C401; DM00013 P23274 18-306: P118-C401 PUTATIVE G-PROTEIN COUPLED RECEPTOR, RALC: PD170483: V345-F403 G-protein coupled receptor: BL00237A: Q187-P226; BL00237D: P380-C396 Olfactory receptor signature PR00245A: M156-K177; PR00245B: S274-D288; PR00245C: F335-V350 EDG1 orphan receptor signature (a G-coupled protein receptor): PR00642D: I145-F159 Transmembrane domain: W123-E149; M156-M175; I299-I318 7-transmembrane receptor (rhodopsin family; 7 <sup>tm</sup> 1): G138-Y388 G-Protein Receptor motif: V207-I223	HMME HMME-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO BLAST-DOMO BLAST-PRODROM BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PRINTS HMME HMME-PFAM MOTIFS
8	7474839CD1	408	S6, T34, Y157, S164, S205, S285, S326, S327	N139		

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
9	7474872CB1	2444	1-82, 2197-2269, 921-1315	70812700V1	1879	2444
				71703018V1	467	1076
				6251915H1 (LUNPTUT02)	1845	2443
				71691712V1	3	462
				71702908V1	1078	1801
				71563743V1	1566	2343
10	6575963CB1	1014	1-101, 515-1014	71703130V1	1011	1763
				71566266V1	247	738
				GNN.g7528005_000007_004	1	1014
11	7474846CB1	1083	908-1006, 1-159, 249-831	4609810F6 (BRSTTUT15)	1	274
				GNN.g7276252_000047_002	43	1083
12	1258785CB1	1740	1-1013	3271244H1 (BRAINOT20)	668	892
				2378395F6 (ISLTNOT01)	865	1366
				1345270F6 (PROSNOT11)	216	821
				SBAA00212F1	1426	1740
				SBAA02650F1	970	1429
				g5663306	1	365
				SBAA02409F1	1283	1740
				2653626H1 (THYMNOT04)	131	371
				1676788T6 (STOMFET01)	2317	2965
				70706752V1	1401	2057
13	1874944CB1	3002	1-547, 2980-3002, 651-1921	70708707V1	778	1445
				2954255F6 (KIDNFET01)	2589	3002
				7626751J1 (GBLADIE01)	646	1152
				2656844T6 (LUNGTUT09)	2019	2946
				70704786V1	1441	2115
				6805278J1 (SKIRNOR01)	1	728
14	7475270CB1	965	754-965, 1-717	2939091T6 (THYMFET02)	632	965
				GNN.g7239420_000072_004	1	816
15	55000189CB1	1617	1-214, 1280-1617, 347-732	GS.55000189.fasta	688	1617
				71689627V1	1	735
16	7474839CB1	1227	691-1227, 1-631	GNN.g7235307_000007_004	1	1227

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
9	7474872CB1	MENITUT03
10	6575963CB1	COLHTUS02
11	7474846CB1	BRSTTUT15
12	1258785CB1	BRAITUT08
13	1874944CB1	KIDNFET01
14	7475270CB1	THYMFET02
15	55000189CB1	LIVRDIR01

Table 6

Library	Vector	Library Description
BRAITUT08	pINCY	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 47-year-old Caucasian male during excision of cerebral meningeal tissue. Pathology indicated grade 4 fibrillary astrocytoma with focal tumoral radionecrosis. Patient history included cerebrovascular disease, deficiency anemia, hyperlipidemia, epilepsy, and tobacco use. Family history included cerebrovascular disease and a malignant prostate neoplasm.
BRSTTUT15	pINCY	Library was constructed using RNA isolated from breast tumor tissue removed from a 46-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 3, nuclear grade 2 adenocarcinoma, ductal type. An intraductal carcinoma component, non-comedo, comprised approximately 50% of the neoplasm, including the lactiferous ducts. Angiolymphatic involvement was present. Metastatic adenocarcinoma was present in 7 of 10 axillary lymph nodes. The largest nodal metastasis measured 3 cm, and focal extracapsular extension was identified. Family history included atherosclerotic coronary artery disease, type II diabetes, cerebrovascular disease, and depressive disorder.
COLHTUS02	pINCY	This subtracted colon tumor tissue library was constructed using 4.24 million clones from a colon tumor library and was subjected to two rounds of subtraction hybridization with 4.04 million clones from an ascending/transverse colon tissue library. The starting library for subtraction was constructed using RNA isolated from colon tumor tissue removed from the hepatic flexure of a 55-year-old Caucasian male during right hemicolectomy, incidental appendectomy, and permanent colostomy. Pathology indicated invasive grade 3 adenocarcinoma that formed a circumferential mass in the ascending colon, located 10.5 cm from the distal resection margin. The tumor infiltrated through the muscularis propria into the pericolononic adipose tissue to within 0.4 cm of the radial fat margin. Patient history included benign hypertension, anxiety, abnormal blood chemistry, blepharitis, heart block, osteoporosis, and hyperplasia of prostate. Family history included prostate cancer, acute myocardial infarction, stroke, and atherosclerotic coronary artery disease. The hybridization probe for subtraction was derived from a similarly constructed library using RNA isolated from non-tumorous ascending and transverse colon tissue from the same donor. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR 19 (1991):1954 and Bonaldo, et al., Genome Research 6 (1996):791.

Table 6 (cont.)

Library	Vector	Library Description
KIDNFET01	pINCY	Library was constructed using RNA isolated from kidney tissue removed from a Caucasian female fetus, who died at 17 weeks' gestation from anencephalus.
LIVRDIR01	pINCY	The library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis diagnosed in 1989. Serology was positive for anti-mitochondrial antibody.
MENITUT03	pINCY	Library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
THYMFET02	pINCY	Library was constructed using RNA isolated from thymus tissue removed from a Caucasian female fetus, who died at 17 weeks' gestation from anencephalus.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater



Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score = 3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

Table 8

Tissues	Polynucleotide SEQ ID NO:		
	6	7	8
Breast, Fat, Skin	-	-	+
Muscle, Bone, Synovium, Connective tissue	+	-	+
Pancreas, Liver, Gallbladder	+	+	-
Brain: Amygdala, Thalamus, Hippocampus, Entorhinal cortex, Archacocortex	-	-	+
Brain: Striatum, Caudate nucleus, Putamen, Dentate nucleus, Globus pallidus, Substantia innominata, Ralphhe magnus	-	-	+
Kidney, Fetal colon, Small intestine, Ileum, Esophagus	-	-	-
Fetal heart, Aorta, Coronary artery	-	-	+
Fetal lung, Adult lung	-	-	+
Placenta, Prostate, Uterus	-	-	-
Olfactory bulb	-	-	-

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
  - a) a polypeptide comprising an amino acid sequence selected from the group consisting of  
5 SEQ ID NO:1-8,
  - b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-8,
  - c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and  
10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-8.
- 15 3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID  
20 NO:9-16.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
- 25 7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
9. A method for producing a polypeptide of claim 1, the method comprising:  
30 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and  
b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide selected from the group consisting of:

- 5 of SEQ ID NO:9-16,
- b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16,
  - c) a polynucleotide complementary to a polynucleotide of a),
  - d) a polynucleotide complementary to a polynucleotide of b), and
  - 10 e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

15 13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
- 20 complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25 14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- 30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.

18. A method for treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

10       a) exposing a sample comprising a polypeptide of claim 1 to a compound, and  
       b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

15

21. A method for treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

20       a) exposing a sample comprising a polypeptide of claim 1 to a compound, and  
       b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

25

24. A method for treating a disease or condition associated with overexpression of functional GCREC, comprising administering to a patient in need of such treatment a composition of claim 23.

30

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

      a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and  
35       b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a

compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 5       a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound  
10   with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target  
15   polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- 20    c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- 25    b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- 30    c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of GCREC in a biological sample comprising the steps of:

- 5 a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- 10 a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a F(ab')<sub>2</sub> fragment, or
- e) a humanized antibody.

15 31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of GCREC in a subject, comprising administering to said subject an effective amount of the composition of claim 20 31.

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of GCREC 25 in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

- 30 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, or an immunogenic fragment thereof, under conditions to elicit an antibody response;
- b) isolating antibodies from said animal; and
- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal

antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.

36. An antibody produced by a method of claim 35.

5

37. A composition comprising the antibody of claim 36 and a suitable carrier.

38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

- 10 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, or an immunogenic fragment thereof, under conditions to elicit an antibody response;
- b) isolating antibody producing cells from the animal;
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-
- 15 producing hybridoma cells;
- d) culturing the hybridoma cells; and
- e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.

20 39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

41. The antibody of claim 10, wherein the antibody is produced by screening a Fab

25 expression library.

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

30 43. A method for detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8 in a sample, comprising the steps of:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
- b) detecting specific binding, wherein specific binding indicates the presence of a



polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8 in the sample.

44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8 from a sample, the method comprising:
- incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
  - separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8.

10

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

15

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

20

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

25

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:9.

30

54. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:10.

55. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:11.

56. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:12.

57. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
5 NO:13.

58. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:14.

10 59. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:15.

60. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:16.

<110> INCYTE GENOMICS, INC.  
 PATTERSON, Chandra  
 LU, Dyung Aina M.  
 THORNTON, Michael  
 LU, Yan  
 TRIBOULEY, Catherine M.  
 GRAUL, Richard  
 KHAN, Farrah A.  
 GANDHI, Ameena R.  
 WALIA, Narinder K.  
 NGUYEN, Danniel B.  
 YUE, Henry  
 HAFALIA, April  
 ELLIOTT, Vicki S.  
 LAL, Preeti  
 REDDY, Roopa  
 KALLICK, Deborah A.  
 TANG, Y. Tom  
 AU-YOUNG, Janice

<120> G-PROTEIN COUPLED RECEPTORS

<130> PI-0096 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/205,628; 60/206,222; 60/207,566; 60/208,834; 60/208,861

<151> 2000-05-18; 2000-05-22; 2000-05-25; 2000-06-02; 2000-06-02

<160> 16

<170> PERL Program

<210> 1

<211> 372

<212> PRT

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7474872CD1

<400> 1

Met	Leu	Ala	Asn	Ser	Ser	Ser	Thr	Asn	Ser	Ser	Val	Leu	Pro	Cys
1				5					10					15
Pro	Asp	Tyr	Arg	Pro	Thr	His	Arg	Leu	His	Leu	Val	Val	Tyr	Ser
				20					25					30
Leu	Val	Leu	Ala	Ala	Gly	Leu	Pro	Leu	Asn	Ala	Leu	Ala	Leu	Trp
				35					40					45
Val	Phe	Leu	Arg	Ala	Leu	Arg	Val	His	Ser	Val	Val	Ser	Val	Tyr
				50					55					60
Met	Cys	Asn	Leu	Ala	Ala	Ser	Asp	Leu	Leu	Phe	Thr	Leu	Ser	Leu
				65					70					75
Pro	Val	Arg	Leu	Ser	Tyr	Tyr	Ala	Leu	His	His	Trp	Pro	Phe	Pro
				80					85					90
Asp	Leu	Leu	Cys	Gln	Thr	Thr	Gly	Ala	Ile	Phe	Gln	Met	Asn	Met
				95					100					105
Tyr	Gly	Ser	Cys	Ile	Phe	Leu	Met	Leu	Ile	Asn	Val	Asp	Arg	Tyr
				110					115					120
Ala	Ala	Ile	Val	His	Pro	Leu	Arg	Leu	Arg	His	Leu	Arg	Arg	Pro
				125					130					135
Arg	Val	Ala	Arg	Leu	Leu	Cys	Leu	Gly	Val	Trp	Ala	Leu	Ile	Leu

Val	Phe	Ala	Val	140	Pro	Ala	Ala	Arg	Val	145	His	Arg	Pro	Ser	Arg	Cys	150
Arg	Tyr	Arg	Asp	155	Leu	Glu	Val	Arg	Leu	160	Cys	Phe	Glu	Ser	Phe	Ser	165
Asp	Glu	Leu	Trp	170	Lys	Gly	Arg	Leu	Leu	175	Pro	Leu	Val	Leu	Leu	Ala	180
Glu	Ala	Leu	Gly	185	Phe	Leu	Leu	Pro	Leu	190	Ala	Ala	Val	Val	Tyr	Ser	195
Ser	Gly	Arg	Val	200	Phe	Trp	Thr	Leu	Ala	205	Arg	Pro	Asp	Ala	Thr	Gln	210
Ser	Gln	Arg	Arg	215	Arg	Lys	Thr	Val	Arg	220	Leu	Leu	Leu	Ala	Asn	Leu	225
Val	Ile	Phe	Leu	230	Leu	Cys	Phe	Val	Pro	235	Tyr	Asn	Ser	Thr	Leu	Ala	240
Val	Tyr	Gly	Leu	245	Leu	Arg	Ser	Lys	Leu	250	Val	Ala	Ala	Ser	Val	Pro	255
Ala	Arg	Asp	Arg	260	Val	Arg	Gly	Val	Leu	265	Met	Val	Met	Val	Leu	Leu	270
Ala	Gly	Ala	Asn	275	Cys	Val	Leu	Asp	Pro	280	Leu	Val	Tyr	Tyr	Phe	Ser	285
Ala	Glu	Gly	Phe	290	Arg	Asn	Thr	Leu	Arg	295	Gly	Leu	Gly	Thr	Pro	His	300
Arg	Ala	Arg	Thr	305	Ser	Ala	Thr	Asn	Gly	310	Thr	Arg	Ala	Ala	Leu	Ala	315
Gln	Ser	Glu	Arg	320	Ser	Ala	Val	Thr	Thr	325	Asp	Ala	Thr	Arg	Pro	Asp	330
Ala	Ala	Ser	Gln	335	Gly	Leu	Leu	Arg	Pro	340	Ser	Asp	Ser	His	Ser	Leu	345
Ser	Ser	Phe	Thr	350	Gln	Cys	Pro	Gln	Asp	355	Ser	Ala	Leu				360
				365						370							

&lt;210&gt; 2

&lt;211&gt; 337

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 6575963CD1

&lt;400&gt; 2

Met	Asn	Glu	Pro	Leu	Asp	Tyr	Leu	Ala	Asn	Ala	Ser	Asp	Phe	Pro			
1				5					10					15			
Asp	Tyr	Ala	Ala	Ala	Phe	Gly	Asn	Cys	Thr	Asp	Glu	Asn	Ile	Pro			
				20					25					30			
Leu	Lys	Met	His	Tyr	Leu	Pro	Val	Ile	Tyr	Gly	Ile	Ile	Phe	Leu			
				35					40					45			
Val	Gly	Phe	Pro	Gly	Asn	Ala	Val	Val	Ile	Ser	Thr	Tyr	Ile	Phe			
				50					55					60			
Lys	Met	Arg	Pro	Trp	Lys	Ser	Ser	Thr	Ile	Ile	Met	Leu	Asn	Leu			
				65					70					75			
Ala	Cys	Thr	Asp	Leu	Leu	Tyr	Leu	Thr	Ser	Leu	Pro	Phe	Leu	Ile			
				80					85					90			
His	Tyr	Tyr	Ala	Ser	Gly	Glu	Asn	Trp	Ile	Phe	Gly	Asp	Phe	Met			
				95					100					105			
Cys	Lys	Phe	Ile	Arg	Phe	Ser	Phe	His	Phe	Asn	Leu	Tyr	Ser	Ser			
				110					115					120			
Ile	Leu	Phe	Leu	Thr	Cys	Phe	Ser	Ile	Phe	Arg	Tyr	Cys	Val	Ile			
				125					130					135			
Ile	His	Pro	Met	Ser	Cys	Phe	Ser	Ile	His	Lys	Thr	Arg	Cys	Ala			
				140					145					150			
Val	Val	Ala	Cys	Ala	Val	Val	Trp	Ile	Ile	Ser	Leu	Val	Ala	Val			

Ile Pro Met Thr	155	160	165
Phe Leu Ile Thr Ser	170	175	180
Ser Ala Cys Leu Asp	185	190	195
Lys Trp Tyr Asn Leu	200	205	210
Leu Val Ile Val Thr	215	220	225
Thr His Gly Leu Gln	230	235	240
Arg Leu Thr Ile Leu	245	250	255
Pro Phe His Ile Leu	260	265	270
Ser Ile Ser Cys Ser	275	280	285
Val Ser Arg Pro Leu	290	295	300
Leu Tyr Val Val Ser	305	310	315
Thr Val Arg Cys Lys	320	325	330
Ile Ser Tyr Ser Asn	335		

&lt;210&gt; 3

&lt;211&gt; 346

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7474846CD1

&lt;400&gt; 3

Met Tyr Asn Gly Ser	5	10	15
Gln Val Met Pro Pro	20	25	30
Leu Gly Asn Gly Val	35	40	45
Thr Trp Lys Pro Ser	50	55	60
Asp Phe Leu Leu Met	65	70	75
Leu Arg Arg Arg His	80	85	90
Gly Leu Phe Thr Leu	95	100	105
Leu Thr Val Val Ala	110	115	120
His His Ala Val Asn	125	130	135
Val Cys Thr Leu Trp	140	145	150
Leu Leu Glu Asn His	155	160	165
Glu Ser Phe Ile Met	170	175	180
Phe Gln Leu Glu Phe	185	190	195
Ser Phe Lys Ile Val			

	200		205		210
Arg Gln Ala Arg	Met Lys Lys Ala Thr	Arg Phe Ile Met Val	Val		
	215		220		225
Ala Ile Val Phe	Ile Thr Cys Tyr Leu	Pro Ser Val Ser Ala	Arg		
	230		235		240
Leu Tyr Phe Leu	Trp Thr Val Pro Ser	Ser Ala Cys Asp Pro	Ser		
	245		250		255
Val His Gly Ala	Leu His Ile Thr Leu	Ser Phe Thr Tyr Met	Asn		
	260		265		270
Ser Met Leu Asp	Pro Leu Val Tyr Tyr	Phe Ser Ser Pro Ser	Phe		
	275		280		285
Pro Lys Phe Tyr	Asn Lys Leu Lys Ile	Cys Ser Leu Lys Pro	Lys		
	290		295		300
Gln Pro Gly His	Ser Lys Thr Gln Arg	Pro Glu Glu Met Pro	Ile		
	305		310		315
Ser Asn Leu Gly	Arg Arg Ser Cys Ile	Ser Val Ala Asn Ser	Phe		
	320		325		330
Gln Ser Gln Ser	Asp Gly Gln Trp Asp	Pro His Ile Val Glu	Trp		
	335		340		345
His					

<210> 4  
 <211> 432  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1258785CD1

<400> 4

Met Glu Glu Arg Ala	Phe Val Asn Pro	Phe Pro Asp Tyr Glu	Ala
1	5	10	15
Ala Ala Gly Ala Leu	Leu Ala Ser Gly	Ala Ala Glu Glu Thr	Gly
	20	25	30
Cys Val Arg Pro Pro	Ala Thr Thr Asp	Glu Pro Gly Leu Pro	Phe
	35	40	45
His Gln Asp Gly Lys	Ile Ile His Asn	Phe Ile Arg Arg Ile	Gln
	50	55	60
Thr Lys Ile Lys Asp	Leu Leu Gln Gln	Met Glu Glu Gly Leu	Lys
	65	70	75
Thr Ala Asp Pro His	Asp Cys Ser Ala	Tyr Thr Gly Trp Thr	Gly
	80	85	90
Ile Ala Leu Leu Tyr	Leu Gln Leu Tyr	Arg Val Thr Cys Asp	Gln
	95	100	105
Thr Tyr Leu Leu Arg	Ser Leu Asp Tyr	Val Lys Arg Thr Leu	Arg
	110	115	120
Asn Leu Asn Gly Arg	Arg Val Thr Phe	Leu Cys Gly Asp Ala	Gly
	125	130	135
Pro Leu Ala Val Gly	Ala Val Ile Tyr	His Lys Leu Arg Ser	Asp
	140	145	150
Cys Glu Ser Gln Glu	Cys Val Thr Lys	Leu Leu Gln Leu Gln	Arg
	155	160	165
Ser Val Val Cys Gln	Glu Ser Asp Leu	Pro Asp Glu Leu Leu	Tyr
	170	175	180
Gly Arg Ala Gly Tyr	Leu Tyr Ala Leu	Leu Tyr Leu Asn Thr	Glu
	185	190	195
Ile Gly Pro Gly Thr	Val Cys Glu Ser	Ala Ile Lys Glu Val	Val
	200	205	210
Asn Ala Ile Ile Glu	Ser Gly Lys Thr	Leu Ser Arg Glu Glu	Arg
	215	220	225
Lys Thr Glu Arg Cys	Pro Leu Leu Tyr	Gln Trp His Arg Lys	Gln

	230		235		240
Tyr Val Gly Ala	Ala His Gly Met Ala	Gly Ile Tyr Tyr Met	Leu		
	245		250		255
Met Gln Pro Ala	Ala Lys Val Asp Gln	Glu Thr Leu Thr Glu	Met		
	260		265		270
Val Lys Pro Ser	Ile Asp Tyr Val Arg	His Lys Lys Phe Arg	Ser		
	275		280		285
Gly Asn Tyr Pro	Ser Ser Leu Ser Asn	Glu Thr Asp Arg Leu	Val		
	290		295		300
His Trp Cys His	Gly Ala Pro Gly Val	Ile His Met Leu Met	Gln		
	305		310		315
Ala Tyr Lys Val	Phe Lys Glu Glu Lys	Tyr Leu Lys Glu Ala	Met		
	320		325		330
Glu Cys Ser Asp	Val Ile Trp Gln Arg	Gly Leu Leu Arg Lys	Gly		
	335		340		345
Tyr Gly Ile Cys	His Gly Thr Ala Gly	Asn Gly Tyr Ser Phe	Leu		
	350		355		360
Ser Leu Tyr Arg	Leu Thr Gln Asp Lys	Lys Tyr Leu Tyr Arg	Ala		
	365		370		375
Cys Lys Phe Ala	Glu Trp Cys Leu Asp	Tyr Gly Ala His Gly	Cys		
	380		385		390
Arg Ile Pro Asp	Arg Pro Tyr Ser Leu	Phe Glu Gly Met Ala	Gly		
	395		400		405
Ala Ile His Phe	Leu Ser Asp Val Leu	Gly Pro Glu Thr Ser	Arg		
	410		415		420
Phe Pro Ala Phe	Glu Leu Asp Ser Ser	Lys Arg Asp			
	425		430		

&lt;210&gt; 5

&lt;211&gt; 240

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1874944CD1

&lt;400&gt; 5

Met Pro Val Leu Leu	His Tyr Phe Phe	Leu Ser Ala Phe Ala	Trp
1	5	10	15
Met Leu Val Glu Gly	Leu His Leu Tyr	Ser Met Val Ile Lys	Val
	20	25	30
Phe Gly Ser Glu Asp	Ser Lys His Arg	Tyr Tyr Gly Met	Gly
	35	40	45
Trp Gly Phe Pro Leu	Leu Ile Cys Ile	Ile Ser Leu Ser Phe	Ala
	50	55	60
Met Asp Ser Tyr Gly	Thr Ser Asn Asn	Cys Trp Leu Ser Leu	Ala
	65	70	75
Ser Gly Ala Ile Trp	Ala Phe Val Ala	Pro Ala Leu Phe Val	Ile
	80	85	90
Val Val Asn Ile Gly	Ile Leu Ile Ala	Val Thr Arg Val Ile	Ser
	95	100	105
Gln Ile Ser Ala Asp	Asn Tyr Lys Ile	His Gly Asp Pro Ser	Ala
	110	115	120
Phe Lys Leu Thr Ala	Lys Ala Val Ala	Val Leu Leu Pro Ile	Leu
	125	130	135
Gly Thr Ser Trp Val	Phe Gly Val Leu	Ala Val Asn Gly Cys	Ala
	140	145	150
Val Val Phe Gln Tyr	Met Phe Ala Thr	Leu Asn Ser Leu Gln	Gly
	155	160	165
Leu Phe Ile Phe Leu	Phe His Cys Leu	Leu Asn Ser Glu Val	Arg
	170	175	180
Ala Ala Phe Lys His	Lys Thr Lys Val	Trp Ser Leu Thr Ser	Ser

	185		190		195
Ser Ala Arg Thr	Ser Asn Ala Lys Pro	Phe His Ser Asp Leu	Met		
	200		205		210
Asn Gly Thr Arg	Pro Gly Met Ala Ser	Thr Lys Leu Ser Pro	Trp		
	215		220		225
Asp Lys Ser Ser	His Ser Ala His Arg	Val Asp Leu Ser Ala	Val		
	230		235		240

&lt;210&gt; 6

&lt;211&gt; 271

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7475270CD1

&lt;400&gt; 6

Met Gly Arg Trp Val	Asn Gln Ser Tyr Thr	Asp Gly Phe Phe Leu	
1	5	10	15
Leu Gly Ile Phe Ser	His Ser Gln Thr Asp	Leu Val Leu Phe Ser	
	20	25	30
Ala Val Met Val Val	Phe Thr Val Ala Leu	Cys Gly Asn Val Leu	
	35	40	45
Leu Ile Phe Leu Ile	Tyr Leu Asp Ala Gly	Leu His Thr Pro Met	
	50	55	60
Tyr Phe Phe Leu Ser	Gln Leu Ser Leu Met	Asp Leu Met Leu Val	
	65	70	75
Cys Asn Ile Val Pro	Lys Met Ala Ala Asn	Phe Leu Ser Gly Arg	
	80	85	90
Lys Ser Ile Ser Phe	Val Gly Cys Gly Ile	Gln Ile Gly Phe Phe	
	95	100	105
Val Ser Leu Val Gly	Ser Glu Gly Leu Leu	Leu Gly Leu Met Ala	
	110	115	120
Tyr Asp Arg Tyr Val	Ala Val Ser His Pro	Leu His Tyr Pro Ile	
	125	130	135
Leu Met Asn Gln Arg	Val Cys Leu Gln Ile	Thr Gly Ser Ser Trp	
	140	145	150
Ala Phe Gly Ile Ile	Asp Gly Val Ile Gln	Met Val Ala Ala Met	
	155	160	165
Gly Leu Pro Tyr Cys	Gly Ser Arg Ser Val	Asp His Phe Phe Trp	
	170	175	180
Ala Val Leu Arg Ile	Arg Ser Ala Gln Ala	Trp Lys Lys Ala Leu	
	185	190	195
Ala Thr Cys Ser Ser	His Leu Thr Ala Val	Thr Leu Phe Tyr Gly	
	200	205	210
Ala Ala Met Phe Met	Tyr Leu Arg Pro Arg	Arg Tyr Arg Ala Pro	
	215	220	225
Ser His Asp Lys Val	Ala Ser Ile Phe Tyr	Thr Val Leu Thr Pro	
	230	235	240
Met Leu Asn Pro Leu	Ile Tyr Ser Leu Arg	Asn Gly Glu Val Met	
	245	250	255
Gly Ala Leu Arg Lys	Gly Leu Asp Arg Cys	Arg Ile Gly Ser Gln	
	260	265	270

His

&lt;210&gt; 7

&lt;211&gt; 276

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens



&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 55000189CD1

&lt;400&gt; 7

Met	Arg	Arg	Lys	Asn	Leu	Thr	Glu	Val	Thr	Glu	Phe	Val	Phe	Leu
1				5					10					15
Gly	Phe	Ser	Arg	Phe	His	Lys	His	His	Ile	Thr	Leu	Phe	Val	Val
				20					25					30
Phe	Leu	Ile	Leu	Tyr	Thr	Leu	Thr	Val	Ala	Gly	Asn	Ala	Ile	Ile
				35					40					45
Met	Thr	Ile	Ile	Cys	Ile	Asp	Arg	His	Leu	His	Thr	Pro	Met	Tyr
				50					55					60
Phe	Phe	Leu	Ser	Met	Leu	Ala	Ser	Ser	Lys	Thr	Val	Tyr	Thr	Leu
				65					70					75
Phe	Ile	Ile	Pro	Gln	Met	Leu	Ser	Ser	Phe	Val	Thr	Gln	Thr	Gln
				80					85					90
Pro	Ile	Ser	Leu	Ala	Gly	Cys	Thr	Thr	Gln	Thr	Phe	Phe	Phe	Val
				95					100					105
Thr	Leu	Ala	Ile	Asn	Asn	Cys	Phe	Leu	Leu	Thr	Val	Met	Gly	Tyr
				110					115					120
Asp	His	Tyr	Met	Ala	Ile	Cys	Asn	Pro	Leu	Arg	Tyr	Arg	Val	Ile
				125					130					135
Thr	Ser	Lys	Lys	Val	Cys	Val	Gln	Leu	Val	Cys	Gly	Ala	Phe	Ser
				140					145					150
Ile	Gly	Leu	Ala	Met	Ala	Ala	Val	Gln	Val	Thr	Ser	Ile	Phe	Thr
				155					160					165
Leu	Pro	Phe	Cys	His	Thr	Val	Val	Gly	His	Phe	Phe	Cys	Asp	Ile
				170					175					180
Leu	Pro	Val	Met	Lys	Leu	Ser	Cys	Ile	Asn	Thr	Thr	Ile	Asn	Glu
				185					190					195
Ile	Ile	Asn	Phe	Val	Val	Arg	Leu	Phe	Val	Ile	Leu	Val	Pro	Met
				200					205					210
Gly	Leu	Val	Phe	Ile	Ser	Tyr	Val	Leu	Ile	Ile	Ser	Thr	Val	Leu
				215					220					225
Lys	Ile	Ala	Ser	Ala	Glu	Gly	Trp	Lys	Lys	Thr	Phe	Ala	Thr	Cys
				230					235					240
Ala	Phe	His	Leu	Thr	Val	Val	Ile	Val	His	Tyr	Gly	Cys	Ala	Ser
				245					250					255
Ile	Ala	Tyr	Leu	Met	Pro	Lys	Ser	Glu	Asn	Ser	Ile	Glu	Gln	Asp
				260					265					270
Leu	Leu	Leu	Ser	Val	Thr									
				275										

&lt;210&gt; 8

&lt;211&gt; 408

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7474839CD1

&lt;400&gt; 8

Met	Lys	Lys	Arg	Glu	Ser	Leu	Thr	Gln	Leu	Arg	Ser	Pro	Trp	Val
1				5					10					15
Val	Ser	Val	Phe	Gly	Ala	Leu	Ile	Thr	Val	Ala	Arg	Phe	Leu	Asp
				20					25					30
Leu	Val	Pro	Thr	Gln	Arg	Asn	Phe	Phe	Lys	Pro	Val	Arg	Pro	Val
				35					40					45
Pro	Ser	Phe	Ala	Tyr	Pro	Leu	Ser	Gln	Asp	Arg	Thr	Pro	Gln	Phe
				50					55					60
Leu	Pro	Pro	Thr	Leu	His	Leu	Ser	Lys	Ala	Arg	Gly	Ile	Thr	Leu

Pro Gly Lys Lys Tyr	65	Pro Gly Phe Cys Met	70	Gln Lys Pro Gln Leu	75
Leu Val Pro Ile Ile	80	Ala Thr Ser Asn Gly	85	Asn Leu Val His Ala	90
Ala Tyr Phe Leu Leu	95	Val Gly Ile Pro Gly	100	Leu Gly Pro Thr Ile	105
His Phe Trp Leu Ala	110	Phe Pro Leu Cys Phe	115	Met Tyr Ala Leu Ala	120
Thr Leu Gly Asn Leu	125	Thr Ile Val Leu Ile	130	Ile Arg Val Glu Arg	135
Arg Leu His Glu Pro	140	Met Tyr Leu Phe Leu	145	Ala Met Leu Ser Thr	150
Ile Asp Leu Val Leu	155	Ser Ser Ile Thr Met	160	Pro Lys Met Ala Ser	165
Leu Phe Leu Met Gly	170	Ile Gln Glu Ile Glu	175	Phe Asn Ile Cys Leu	180
Ala Gln Met Phe Leu	185	Ile His Ala Leu Ser	190	Ala Val Glu Ser Ala	195
Val Leu Leu Ala Met	200	Ala Phe Asp Arg Phe	205	Val Ala Ile Cys His	210
Pro Leu Arg His Ala	215	Ser Val Leu Thr Gly	220	Cys Thr Val Ala Lys	225
Ile Gly Leu Ser Ala	230	Leu Thr Arg Gly Phe	235	Val Phe Phe Phe Pro	240
Leu Pro Phe Ile Leu	245	Lys Trp Leu Ser Tyr	250	Cys Gln Thr His Thr	255
Val Thr His Ser Phe	260	Cys Leu His Gln Asp	265	Ile Met Lys Leu Ser	270
Cys Thr Asp Thr Arg	275	Val Asn Val Val Tyr	280	Gly Leu Phe Ile Ile	285
Leu Ser Val Met Gly	290	Val Asp Ser Leu Phe	295	Ile Gly Phe Ser Tyr	300
Ile Leu Ile Leu Trp	305	Ala Val Leu Glu Leu	310	Ser Ser Arg Arg Ala	315
Ala Leu Lys Ala Phe	320	Asn Thr Cys Ile Ser	325	His Leu Cys Ala Val	330
Leu Val Phe Tyr Val	335	Pro Leu Ile Gly Leu	340	Ser Val Val His Arg	345
Leu Gly Gly Pro Thr	350	Ser Leu Leu His Val	355	Val Met Ala Asn Thr	360
Tyr Leu Leu Leu Pro	365	Pro Val Val Asn Pro	370	Leu Val Tyr Gly Ala	375
Lys Thr Lys Glu Ile	380	Cys Ser Arg Val Leu	385	Cys Met Phe Ser Gln	390
Gly Gly Lys	395		400		405

&lt;210&gt; 9

&lt;211&gt; 2444

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7474872CB1

&lt;400&gt; 9

```

cgtgagatct gctgaggtgg gtgtgtcccc tcccgccccg ggagcaggtc ctaccagccc 60
agcccagccc agcccagccc agagcaggca gcggaagcca gcttggggca ggcgagagca 120
acacggagca caggtctctg ctgctgatga agctgtgacc aaacgcaccc aacccttggc 180
agccatctgt ccctgcagcc atagcccaca ttcccatgac ctccctctgc ttgttttggg 240
accatgtctg tacagcctct agggcccagc cccggaggtg aatgccatgc catgattctg 300

```

```

gtgtgtctcca tggcatcccc agcctagctc ccaatcccac tttggcacga tgttagccaa 360
cagctcctca accaacagtt ctgtttctcc gtgtcctgac taccgacctt cccaccgctt 420
gcacttggtg gtctacagct tgggtgctggc tgccgggctc cccctcaacg cgctagccct 480
ctgggtcttc ctgctgctgc tgcgctgca ctggtggtg agcgtgtaca tgtgtaacct 540
ggcggccagg gacctgtctc tcacctctc gctgcccgtt cgtctctcct actacgact 600
gcaccactgg ccttccccg acctcctgtg ccagacgacg ggcgccatct tccagatgaa 660
catgtacggc agctgcatct tctgatgct catcaacgtg gaccgctacg ccgcatcgt 720
gcacccgctg cgactgcgcc acctgcggcg gccccgctg gcgcggtgc tctgctggg 780
cgtgtgggcg ctcatcctgg tgtttgcct gcccgcggc cgctgcaca ggccctcgcg 840
ttgcgctac cgggacctg aggtgcgct atgcttcgag agcttcagcg acgagctgtg 900
gaaaggcctg ctgctgccc tctgtctgt ggcgaggcg ctgggcttcc tctgccccct 960
ggcggcggtg gtctactcgt cgggcccagt cttctggacg ctggcgcgcc ccgacgccac 1020
gcagagccag cggcgggcga agaccgtgc cctcctgctg gctaacctcg tcatcttct 1080
gctgtgcttc gtgccctaca acagcacgt ggcggtctac gggctgctgc ggagcaagct 1140
ggtggcgggc agcgtgcctg cccgcgatcg cgtgcgcggg gtgctgatgg tgatgggtgt 1200
gctggcgggc gccaaactgc tgcctggacc gctggtgtac tactttagcg ccgagggctt 1260
ccgcaacacc ctgctgccc tgggcactcc gcaccgggccc aggacctcg ccaccaacgg 1320
gacgcgggcg gcgctcgcg aatccgaaag gtccgcccgc accaccgacg ccaccaggcc 1380
ggatgcggcc agtcaggggc tgcctcgacc ctccgactcc cactctctgt cttccttcac 1440
acagtgtccc caggattccg cctctgaac acacatgcca ttgcgctgtc cgtgcccgc 1500
tcccaacgcc tctcgttctg ggaggcttac aggtgtaca cacaagaagg tgggctgggc 1560
acttggaact ttgggtggca attccagctt agcaacgcag aagagtacaa agtgtggaag 1620
ccagggccca gggaaggcag tgcgtcttct aatggcttct ttaaactgtg agcacgcaga 1680
gcacccttc tccagcggtg ggaagtgatg cagagagccc acccgtgcag agggcagaag 1740
aggacgaaat gcctttgggt gggcagggca ttaaactgct aaaagctggt tagatggaac 1800
agaaaatggg cattctggat ctaaaccgcc acaggggcct gagagctgaa gagaccagg 1860
tttgggtggc aaagctactg agatgcctgt tcatctgctg acttctgtct aggtcatgg 1920
atgccacctt ctttcatttc ggcctaggct tcccctgctc accactgagg cctaatacaa 1980
gagttcctat ggacagaact acattcttcc tcgcatagtg acttgtaga atttagactt 2040
ggcatccagc atgggatagt tggggcaagg caaaactaac ttagagtctt cccctcaaca 2100
acatccaagt ccaaaccctt ttaggttat cctttcttcc atcacatccc cttttccagg 2160
cctcctccat ttaggtcct taatattctt tcttttctc tctctctcgt ttctctctt 2220
tctctctctt cctctctctt ctctctctt ctctctctcc cgctctctcc tttgtccaga 2280
gtaaggataa aattctttct actaaagcac tgggttctca actttttggt ctcagacccc 2340
actcttagaa attgaggatc tcaaagagct ttgcttatat tttgttctt tgatacttac 2400
catactagaa attaaagcga atacattttt aaaataaaaa aaaa 2444

```

&lt;210&gt; 10

&lt;211&gt; 1014

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 6575963CB1

&lt;400&gt; 10

```

atgaatgagc cactagacta ttagcaaat gcttctgatt tccccgatta tgcagctgct 60
tttggaaatt gcactgatga aaacatccca ctcaagatgc actacctccc tgttatattat 120
ggcattatct tctcgtggg atttccaggc aatgcagtag tgatatccac ttacattttc 180
aaaatgagac cttggaagag cagcaccatc attatgctga acctggcctg cacagatctg 240
ctgtatctga ccagcctccc cttcctgatt cactactatg ccagtggcga aaactggatc 300
tttggagatt tcatgtgtaa gtttatccgc ttcagcttcc atttcaacct gtatagcagc 360
atcctcttcc tcacctgttt cagcatcttc cgctactgtg tgatcattca ccaatgagc 420
tgcttttcca ttcacaaaac tcgatgtgca gttgtagcct gtgctgtggg gtggatcatt 480
tactggtag ctgtcattcc gatgaccttc ttgatcacat caaccaacag gaccaacaga 540
tcagcctgtc tcgacctcac cagtccggat gaactcaata ctattaagtg gtacaacctt 600
attttgactg caactacttt ctgcctcccc ttggtgatag tgacactttg ctataccacg 660
attatccaca ctctgaccca tggactgcaa actgacagct gccttaagca gaaagcagca 720
agggtacaac tctgtact ccttgcaatt tacgtatgtt tttaccctt ccatactctg 780
agggtcattc ggatcgaatc tcgctgtctt tcaatcagtt gttccattga gaatcagatc 840
catgaagctt acatcgtttc tagaccatta gctgctctga acacctttgg taacctgtta 900
ctatatgtgg tggtcagcga caactttcag caggctgtct gctcaacagt gagatgcaaa 960

```

gtaagcggga accttgagca agcaaagaaa attagttact caaacaaccc ttga 1014

<210> 11  
 <211> 1083  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7474846CB1

<400> 11  
 ccctgggctg ctctgcaccc ggacacttgc tctgtcccg ccatgtacaa cgggtcgtgc 60  
 tgccgcatcg agggggacac catctcccag gtgatgccgc cgctgctcat tgtggccttt 120  
 gtgctgggag cactaggcaa tggggctgcc ctgtgtggtt tctgcttcca catgaagacc 180  
 tggaagccca gcactgttta ccttttcaat ttggccgtgg ctgatttcct ccttatgac 240  
 tgccctgcctt ttccggacaga ctattacctc agacgtagac actgggcttt tggggacatt 300  
 ccctgccgag tggggctctt cacgttggcc atgaacaggg ccgggagcat cgtgttcctt 360  
 acggtgggtg ctgaggacag gtatttcaaa gtgttccacc cccaccacgc ggtgaacact 420  
 atctccaccc ggggtggcggc tggcatcgct tgcaccctgt gggccctggg catcctggga 480  
 acagtgtatc ttttgctgga gaacctctc tgcgtgcaag agacggcgt ctctgtgag 540  
 agcttcatca tggagtcggc caatggctgg catgacatca tgttcagct ggagttcttt 600  
 atgccctcgc gcatcatctt attttgcctc ttcaagattg tttggagcct gaggcggagg 660  
 cagcagctgg ccagacaggc tcggatgaag aaggcgaccc ggttcatcat ggtggtggca 720  
 attgtgttca tcacatgcta cctgccacgc gtgtctgcta gactctattt cctctggacg 780  
 gtgccctcga gtgcctgcga tccctctgtc catggggccc tgcacataac cctcagcttc 840  
 acctacatga acagcatgct ggatccccctg gtgtattatt tttcaagccc ctcccttccc 900  
 aaattctcaa acaagctcaa aatctgcagt ctgaaaccca agcagccagg acactcaaaa 960  
 acacaaaggc cgggaagagat gccaatctcg aacctcggtc gcaggagtgt catcagttg 1020  
 gcaaatagtt tccaaagcca gtctgatggg caatgggatc cccacattgt tgagtggcac 1080  
 tga 1083

<210> 12  
 <211> 1740  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1258785CB1

<400> 12  
 ggcagtgcac gctcagacgc cccgctcctc ccgccagcgc gcggcctcgc tcctcctaga 60  
 ggacgctctc tgccgaggcc ctccgaggag gcggcgccgc ggcgagctgc agcgccggga 120  
 caggaggttt gtccccgccc gcgcgccgta ccgcggcgga gatgggagag accatgtcaa 180  
 agaggctgaa gctccacctg ggaggggagg cagaatgcca ggaacgggcg ttcgtcaacc 240  
 ccttcccga ctacgaggcc gccgcggggc cgctgctcgc ctccggagcg gccgaagaga 300  
 caggctgtgt tcgtcccccg gcgaccacgg atgagcccg cctccctttt catcaggacg 360  
 ggaagatcat tcataatttc ataagacgga tccagaccaa aattaaagat cttctgcagc 420  
 aaatggaaga agggctgaag acggctgac cccatgactg ctctgcttat actggctgga 480  
 caggcatagc ccttttgtac ctgcagttgt accgggtcac atgtgaccaa acctacctgc 540  
 tccgatccct ggattacgta aaaagaacac ttccggaatc gaatggccgc agggtcacct 600  
 tcctctgttg ggtgctggc cccctggctg ttggagctgt gatttatcac aaactcagaa 660  
 gtgactgtga gtcccaggaa tgtgtcacia aacttttgca gctccagaga tcggttgtct 720  
 gccagaatc agaccttcct gatgagctgc tttatggacg ggcaggttat ctgtatgcct 780  
 tactgtacct gaacacagag ataggtccag gcaccgtgtg tgagtacgct attaaagagg 840  
 tagtcaatgc tattattgaa tcgggtgaaga ctttgtcaag ggaagaaaga aaaacggagc 900  
 gctgcccgtt gttgtaccag tggcaccgga agcagtagct tggagcagcc catggcatgg 960  
 ctggaattta ctatagtta atgcagccgc gccacaaaagt ggaccaagaa accttgacag 1020  
 aaatggtgaa acccagtatt gattatgtgc gccacaaaaa attccgatct gggaattacc 1080  
 catcatcatt aagcaatgaa acagaccggc tgggtgactg gtgccacggc gccccggggg 1140  
 tcatccacat gctcatgcag gcgtacaagg tctttaagga ggagaagtac ttgaaagagg 1200  
 ccatggagtg tagcgatgtg atttggcagc gaggtttgct gcggaagggc tacgggatat 1260

gccatgggac	tgctggcaac	ggctattcct	tcctgtccct	ttaccgtctc	acgcaggata	1320
agaagtacct	ctaccgagct	tgcaagtttg	cagagtgggtg	tctagattac	ggagcacacg	1380
ggtgccgcat	tcctgacaga	ccctattcgc	tctttgaagg	catggctggc	gctattcact	1440
ttctctctga	tgtcctggga	ccagagacat	cacggtttcc	agcatttgaa	cttgactctt	1500
cgaagaggga	ttaaaagggtg	caaaaagaca	actaaaatac	ccatttggac	caaaagccgc	1560
cagattgctt	agtgcctgac	acagaaacaa	ctgggaatcc	tgaaagagaa	gcagacaccg	1620
tcacaggccc	ctctgggttag	actagcatga	gtgaccgaag	ccatccatca	acattttcta	1680
acagcaccct	catcaatata	aaatatgact	tcttcacata	cagaaaaaaa	aaaaaaaaaa	1740

&lt;210&gt; 13

&lt;211&gt; 3002

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1874944CB1

&lt;400&gt; 13

gcccagttct	ggtgtcdata	gtttccgctt	tgagccgggc	acgacccccct	gccaaagtgat	60
gcccgtgtct	ttacactact	tcttcctgag	tgccttcgca	tggatgctgg	tggagggtct	120
gcacctctac	agcatggtga	tcaaggtctt	tgggtcggag	gacagcaagc	accgttacta	180
ctatgggatg	ggatgggggt	ttcctcttct	gatctgcata	atttactgt	catttgccat	240
ggacagttac	ggaacaagca	acaattgctg	gctgtcggtg	gagagtggcg	ccatctgggc	300
ctttgtagcc	cctgccccgt	ttgtcatcgt	ctacaagatc	ggcatcctca	tcgctgtgac	360
cagagtcata	tcacagatca	gcgccgacaa	ctacaagatc	catggagacc	ccagtgcctt	420
caagttgacg	gccaaggcag	tggcctgtgt	gctgccccatc	ctgggtacct	cgtgggtctt	480
tggcgtgtct	gctgtcaacg	gttgtgtgtg	gggtttccag	tacatgtttg	ccacgctcaa	540
ctccctgcag	ggactgttca	tattcctctt	tcattgtctc	ctgaattcag	aggtgagagc	600
cgccttcaag	cacaaaacca	aggtctgggtc	gctcacgagc	agctccgccc	gcacctccaa	660
cgcgaagccc	ttccactcgg	acctcatgaa	tgggaccccg	ccaggcatgg	cccccaccaa	720
gctcagccct	tgggacaaga	gcagccactc	tggcccaccgc	gtcgacctgt	cagccgtgtg	780
agccgggagg	ctggccaacca	ggccaggctg	cgtctcagaac	acaccccccc	aaacagaatg	840
aaatgccccca	cctttgcccc	tggaccctct	ccttgtgtgt	gtctggacat	gggtgtttgtg	900
gccccgagac	agctgtcctc	ccctgtgact	ctgggtgtcg	gagcacactg	ctcagcccgag	960
cagcctgatg	cccaggccag	cgtgggcccc	cctgccttgc	atccaccctg	gggtgtagtgt	1020
acttctctcg	gggattccca	ggacacagtg	gcctgactgt	gatgggtgcc	ttgagcctcc	1080
cttcatcact	cagcatcaga	cccagcgagg	ccaggacact	cggggccggt	cccgagcac	1140
caggagggga	tgttcagcct	ctgtgccttg	gtggggcctg	gggactcagg	gccaaagagg	1200
tgggttcaggt	ccccacgcac	cctcagtcag	gcgcaggcag	ctgggggtgt	gtgggggaaga	1260
gcatgctggag	tccccagtg	ctgaatccac	tgagtgggtga	gttccccaca	gccggcgcta	1320
gccgtgggtgt	gtgtctctgt	aggtgggtgcc	ggcgtggggc	aacctgtgct	gtgtcatcag	1380
ttggggggccc	ctgcccgaagc	cgagctcgag	ccgtgggcgg	gagtcgttga	ctctccaggt	1440
gagggcgacc	cctctggccc	gtccttgcgg	gggtccccctc	tgctcacgtg	aagagccgct	1500
ctgggccttg	aggctgcctg	atgggtgcctg	tgcttggggg	agcttctcgg	ccatccgctg	1560
tgagttttgc	ctctttggac	cccaattcgg	ccttaagatg	ccctcctccc	tcgtgtgcca	1620
gcctccttgg	ttgttcttgg	gccacaggag	ctggccgtgt	ccccgcagtg	cctggtgtcc	1680
aggtggaaag	tggagggcat	tttccagggc	actgctttcc	ccagaggctt	cctcatggct	1740
cacaggcact	ctacgaagtt	tctaattgggc	agaccacgcg	gcaggtagca	cagtgcgctc	1800
cgtctgggtca	ccatgagacc	gacctgcgct	gagtccccac	tgacctggag	agggaggggt	1860
ggtgacagcc	gtgtcttctg	tgttgaggga	aatttatgga	ctcagactca	gccccagagg	1920
agatgggata	attgttatgg	acccatgtgt	gggcatgata	ctgtggaaca	caggttttggg	1980
atcatagatg	tgaattaaga	caccaccgag	atacgggctg	tgaggttcat	actgtgtctga	2040
tagcactcgt	ggtgtctgtg	aaatgtgggt	aagacattca	aacctgggtt	tgatactgga	2100
aactcttct	ttaaaactgt	gaccatgatt	tcattcagcc	cctccacacc	cctatgtctg	2160
ccttgtttta	gagtgaagtt	tctatggagc	ctgtggccct	tttgcagccc	acctggtggc	2220
ttcttaatgt	aactcttccc	ctggctgcct	ggagtggacc	actcatctgc	aggcctctcc	2280
tgcatggggga	gggttaggcag	ggagcagcat	gtctgcaggg	gtgaaccttt	gctcttctgt	2340
caggcgaggc	ccaggctgca	ccagccacct	gccacatggt	gacagtgcca	cgggcccctgc	2400
gtatggcccc	tgcaaccgtg	ctctggcggg	cacacctggc	tgtctcaggg	caaggccgct	2460
gttcagtgaa	gagtcctcatg	tttagtatgg	actaaagtcc	catgttttagc	catgccccca	2520
gtctcccgtg	accccagaaa	ccaggtcaca	tggaccacag	tgccagatcc	tcctcacgcc	2580
ggtgagcacc	tagaagttag	aacactgtat	tcctacaatg	tacacttgga	tattttctcct	2640

```

tatttagttt ctagtgaaac aaatcaagta aggaactatc ttttagtttag atggaattat 2700
ttgttttttaa ttgttgccgt attcatctat atagctaata tttcaagata agtaatgaac 2760
aaaacctgtc taaacctttt gtttccaatg aatgaaagtc atgcacttta tttataggct 2820
ctatgttttg gcttctgcag tacttttatt atctatacat aatttggcca aaaataagaa 2880
attggaaaga atgaaatgtt tagtttatag tagaagaaag atgatgacac taagtgtgta 2940
aaatatgttg tgatttttat gaaataaact catgtcctga aaaaaaaaaa aaaaaaaaaa 3000
aa 3002

```

&lt;210&gt; 14

&lt;211&gt; 965

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7475270CB1

&lt;400&gt; 14

```

atgggaagat ggggtgaacca gtcctacaca gatggcttct tcctcttggg catcttttcc 60
cacagccaga ctgaccttgt cctcttctct gcagttatgg tggctctcac agtggccctc 120
tgtgggaatg tcctcctcat ctctctcacc tacctggacg ctggacttca caccctcatg 180
tacttcttcc tcagccagct ctccctcatg gacctcatgt tggctctgtaa cattgtgcca 240
aagatggcag ccaacttcct gtctggcagg aagtccatct cctttgtggg ctgtggcata 300
caaattggct tttttgtctc tcttgtggga tctgaggggc tcttgtctgg actcatggct 360
tatgaccgct acgtggccgt tagccaccca ctctactatc ccatcctcat gaatcagagg 420
gtctgtctcc agattactgg gagctcctgg gcctttggga taatagatgg agtgattcag 480
atgggtggcag ccatgggctt accttactgt ggctcaagga gcgtggatca ctttttcttg 540
gctgtgctcc gaatacgctc tgctcaggcc tggaaaaaag ccctggccac ctgctcctcc 600
cacctaacag ctgtcaccct ctctctatgg gcagccatgt tcatgtacct gaggcctagg 660
cgctaccggg cccttagcca tgacaagggt gcctctatct tctacacagt ccttactccc 720
atgctgaacc ccctcattta cagcttgagg aatggggagg tgatgggggc actgaggaag 780
gggctggacc gctgcaggat tggcagccag cactgaaccc cagagtctgg tgcctgctgt 840
gcccttctt gcctgtgtca cattgggaag tcaactcaacc tttgtgagtg tctgtttcca 900
ttcacctgtt gatggtcatt ggatggttta taggttttgg ccatcatgag aaaagccatt 960
atgaa 965

```

&lt;210&gt; 15

&lt;211&gt; 1617

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 55000189CB1

&lt;400&gt; 15

```

gttgcccttt tgtcttgggt cggtgttctg tgtttcgttt tttgtttctc tgttgttgtt 60
tgtgttgtgc gttttctgtt gtgtttttgt gttggtttct ctgttctgcc aggttccagc 120
gcttttgcca ttgtttacgg ccaggctttg gttccgagct tcggattcca tttagtaact 180
gggccgccag ttgtgtgggt ggaattctac atcgtgtcac cagaagctat ccacctatgg 240
ttctaattca gtaagtcaca ctctctcacc cctttttttt gtctcagctg tgtgggcttt 300
cccaggatgg catgcaatgg gacctctgtg ccatgcatat tgtaaaggaa aatgcctccc 360
tccatgcgct acaaaacagc acatttatga tggcactttg aaaagatatg ggctgtggtg 420
tcacatatgg acaattcctt ggccagaagc ttaacagtgc cagcagtgcc agaagattaa 480
gaagacagca aaaacagaaa agggagaaga tgggtgaagta gctatataac atgagcgaga 540
atgctcctga ttacaaagca gagaaattga ctttttttct tagtgttttc tatagtcatt 600
gctctatccc tgttctagaa ttcaagtcac gataagaatt tcttcacgtt gacttccctgc 660
attgctttta gacattgcaa tttaaagaatg cgaagaaaga acctcacaga ggtaacagag 720
tttgttttcc tgggattctc cagattccac aaacatcaca tcactctctt tgtggttttt 780
ctcactctgt acacattaac tgtggctggc aatgccatca tcatgaccat catctgcatt 840
gaccgtcacc tccacactcc catgtacttc ttcttgagca tgctggctag ctcaaagaca 900
gtgtacacac tgttcatcat tccacagatg ctctccagct tcgtaaccca gaccagcca 960
atctccctag caggttgtac caccctaaagc ttcttctttg ttaccttggc catcaacaat 1020

```

```

tgcttcttgc tcacagtgat gggctatgac cactatatgg ccatctgcaa tcccttgaga 1080
tacaggggtca ttacgagcaa gaagggtgtg gtccagctgg tgtgtggagc ctttagcatt 1140
ggcctggcca tggcagctgt ccaggtaaca tccatattta ccttaccttt ttgtcacacg 1200
gtggttggtc atttcttctg tgacatcctc cctgtcatga aactctcctg tattaatacc 1260
actatcaatg agataatcaa ttttgttgtc aggttatttg tcatcctggg ccccatgggt 1320
ctggctcttca tctcctatgt cctcatcatc tccactgtcc tcaagattgc ctcagctgag 1380
ggttggaaga agacctttgc cacctgtgcc ttccacctca ctgtgggtcat tgtccattat 1440
ggctgtgctt ccattgccta cctcatgccc aagtcagaaa actctataga acaagacctc 1500
cttctctcag tgacctaaac catcatcact cccctgtctg accctgttgt ttacagccta 1560
aagaacaagg aggtcaagga tgccctatgc agggccatgg gcagaaacat ttcttaa 1617

```

&lt;210&gt; 16

&lt;211&gt; 1227

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7474839CB1

&lt;400&gt; 16

```

atgaagaagc gagagtcctt gacacagctt cgatccccctt gggtagttag tgtgttcgga 60
gcgctgatca cagtagcccg gttcctggac cttgttcccta cacagaggaa tttcttcaag 120
cctgtgagac ccgttccaag ctttgcctat cctttgtccc aggacaggac tcctcagttc 180
ctgcctccta ctcttcaactt aagcaaggca aggggaatta ctttaccagg aaaaaatac 240
cctggatttt gtatgcagaa gccccagctc ttgggtcccta tcatagccac ttcaaattga 300
aatctgggtc acgcagcata cttccttttg gtgggtatcc ctggcctggg gcctaccata 360
cacttttggc tggctttccc actgtgtttt atgtatgcct tggccaccct gggtaacctg 420
accattgtcc tcatcattcg tgtggagagg cgactgcatg agcccatgta cctcttccctg 480
gccatgcttt ccaactattga cctagtccctc tccctctatca ccatgcccga gatggccagt 540
cttttccctga tgggcatcca ggagatcgag ttcaacattt gcctggccca gatgttccct 600
atccatgctc tgtcagccgt ggagtcagct gtccctgttg ccattggctt tgaccgcttt 660
gtggccattt gccacccatt gcgccatgct tctgtgctga caggggtgtac tgtggccaag 720
attggactat ctgccctgac caggggggtt gtattcttct tcccactgcc cttcatectc 780
aagtggttgt cctactgcca aacacatact gtcacacact ccttctgtct gcaccaagat 840
attatgaagc tgtcctgtac tgacaccagg gtcaatgtgg attggcttct tttatggact cttcatcate 900
ctctcagtca tgggtgtgga ctctctcttc tcggagggca gcaactcaagg ctttcaacac ctgcatctcc 1020
gctgttttgg agctgtcctc cctctatgta cccctcattg ggctctcggg ggtgcatagg 1080
ctgggtgggc ccacctccct cctccatgtg gttatggcta atacctactt gctgctacca 1140
cctgtagtca acccccttgt ctatggagcc aagaccaaaag agatctgttc aagggtccctc 1200
tgtatgttct cacaaggtgg caagtga 1227

```

**THIS PAGE BLANK (USPTO)**